



XXII Geometrical Seminar

May 26 - 31, 2024, Vrnjačka Banja, Serbia

# Book of Abstracts

Editor: Mića Stanković

XXII Geometrical seminar is organized by

Faculty of Sciences and Mathematics, University of Niš and

Faculty of Mathematics, University of Belgrade

in collaboration with

Faculty of Science, University of Kragujevac and

Mathematical Institute of the Serbian Academy of Sciences and Arts



**ISBN 978-86-6275-159-1**

Faculty of Sciences and Mathematics, University of Niš, Serbia  
Faculty of Mathematics, University of Belgrade, Serbia  
Faculty of Science, University of Kragujevac, Serbia  
Mathematical Institute of the Serbian Academy of Sciences and Arts

---



## XXII GEOMETRICAL SEMINAR

### Book of Abstracts

Editor: Mića Stanković

Vrnjačka Banja, 2024.



XXII Geometrical seminar is dedicated to professor  
Stana Nikčević



## Sponsors:



Republic of Serbia

MINISTRY OF SCIENCE,  
TECHNOLOGICAL DEVELOPMENT AND INNOVATION



National Bank of Serbia



**Raiffeisen  
Bank**



**Organizers:** Faculty of Sciences and Mathematics University of Niš, Serbia and Faculty of Mathematics, University of Belgrade, Serbia.

**Co-organizer:** Faculty of Science, University of Kragujevac, Serbia.

**Conference Topics:** Differential Geometry, Topology, Lie Groups, Mathematical Physics, Discrete Geometry, Integrable Systems, Visualization, Teaching Geometry, Geometric Education, as well as other subjects related to the main themes are welcome.

**Governmental sponsors:** Ministry of Science, Technological Development and Innovation.

**International Advisory Committee:**

Dmitri Alekseevsky (Moscow , Russia)  
David Blair (East Lansing, USA)  
Victor Buchstaber (Moscow, Russia)  
Bang-Yen Chen (East Lansing, USA)  
Ryszard Deszcz (Wroclaw, Poland)  
Branko Dragovich (Belgrade, Serbia)  
Anatoly T. Fomenko (Moscow, Russia)  
Graham Hall (Aberdeen, UK)  
Stefan Ivanov (Sofia, Bulgaria)  
Louis Kauffman (Chicago, USA)  
Haizhong Li (Beijing, China)  
Miodrag Mateljević (Belgrade, Serbia)  
Josef Mikeš (Olomouc, Czech Republic)  
Dmitry Millionschikov (Moscow , Russia)  
Svetislav Minčić (Niš, Serbia)  
Andrey Mironov (Novosibirsk, Russia)  
Alexander Mišchenko (Moscow, Russia)  
Emil Molnar (Budapest, Hungary)  
Yuri Nikolayevsky ( Melbourne, Australia)  
Masafumi Okumura (Saitama, Japan)  
Leopold Verstraelen (Leuven, Belgium)  
Andrei Vesnin (Novosibirsk, Russia)  
Iskander Taimanov (Novosibirsk, Russia)  
Alexey Tuzhilin (Moscow, Russia)  
Luc Vrancken (Valenciennes, France)

**Program Committee:**

Vladica Andrejić (Belgrade, Serbia)  
Miroslava Antić (Belgrade, Serbia)  
Mirjana Djorić (Belgrade, Serbia)  
Emilija Nešović (Kragujevac, Serbia)  
Miroslava Petrović-Torgašev (Kragujevac, Serbia)  
Ljiljana Radović (Niš, Serbia)  
Zoran Rakić (Belgrade, Serbia)  
Mića Stanković (Niš, Serbia)  
Ljubica Velimirović (Niš, Serbia)  
Srdjan Vukmirović (Belgrade, Serbia)  
Milan Zlatanović (Niš, Serbia)

**Local Organizing Committee:**

Milica Cvetković (Niš, Serbia)  
Milica Grbović Ćirić (Kragujevac, Serbia)  
Miroslav Maksimović (Kosovska Mitrovica, Serbia)  
Vladislava Milenković (Niš, Serbia)  
Marija Najdanović (Kosovska Mitrovica, Serbia)  
Miloš Petrović (Kruševac, Serbia)  
Mića Stanković (Niš, Serbia)  
Ljubica Velimirović (Niš, Serbia)  
Ana Velimirović (Belgrade, Serbia)  
Nenad Vesić (Belgrade, Serbia)  
Milan Zlatanović (Niš, Serbia)

# Contents

PREFACE . . . . .	1
Sergei Agapov	
- <i>Integrable geodesic flows with rational first integrals</i> . . . . .	3
Dmitri Alekseevsky	
- <i>Hopf Bundle, Listing's Law and Saccades</i> . . . . .	4
Khalifa Al Shaqsi	
- <i>Neutrosophic Touchard Polynomials for Subclass of Analytic Functions</i> . . . . .	5
Miloš Arsenović	
- <i>Convergence theorems for sequences of operator valued functions</i> . . . . .	6
Andronick Arutyunov	
- <i>On a coarse geometric approach to operator algebras</i> . . . . .	7
Muhittin Evren Aydin	
- <i>Pointwise rectifying submanifolds and anti-torqued vector fields</i> . . . . .	8
Vitaly Balashchenko	
- <i>Left-invariant <math>f</math>-structures on low-dimensional solvable Lie groups</i> . . . . .	9
Djordje Baralić	
- <i>The mod <math>p</math> Buchstaber invariant</i> . . . . .	11
Cornelia-Livia Bejan	
- <i>Almost complex manifolds with Norden metrics</i> . . . . .	12
Miguel Brozos-Vázquez	
- <i>Remarks on weighted Einstein manifolds</i> . . . . .	13
Bang-Yen Chen	
- <i>Great antipodal sets and applications</i> . . . . .	14
Milica Cvetković <sup>1</sup> , Ana Mitrović <sup>2</sup>	
- <i>The Willmore energy variations and energy efficient architecture</i> . . . . .	15
Andrijana Dekić	
- <i>Almost Kähler structures on complex hyperbolic space</i> . . . . .	16
Ryszard Deszcz <sup>1</sup> , Małgorzata Głogowska <sup>2</sup> , Jan Jęłowicki <sup>3</sup> , Miroslava Petrović- Torgašev <sup>4</sup> , and Georges Zafindratafa <sup>5</sup>	
- <i>Curvature properties of pseudosymmetry type of some 2-quasi Einstein manifolds</i> . . . . .	17
Ivan Dimitrijević	
- <i>The Schwarzschild-de Sitter Metric of Nonlocal <math>\sqrt{dS}</math> Gravity</i> . . . . .	19
Miloš Djorić	
- <i>Yamabe solitons on submanifolds of Kähler manifolds</i> . . . . .	20
Branko Dragovich	
- <i>Nonlocal de Sitter Gravity <math>\sqrt{dS}</math> and the Dark Side of the Universe</i> . . . . .	21

Vladimir Dragović	
- <i>How to connect two most beautiful geometric theorems?</i>	22
Zdenek Dusek	
- <i>Finsler geodesic orbit metrics</i>	23
Lyudmila Efremova	
- <i>On maps obtained by small perturbations of skew products</i>	24
İrem Küpeli Erken	
- <i>Some Remarks on Quasi-para-Sasakian Manifolds</i>	25
FAN Huijun	
- <i>Spectrum problem of Schrodinger operators related to Landau-Ginzburg models</i>	26
Eduardo Garcia-Rio	
- <i>Critical metrics for quadratic curvature functionals</i>	27
Ilja Gogić	
- <i>Applications of algebraic topology to operator algebras: Homogeneous <math>C^*</math>-algebras</i>	28
Admir Huseini <sup>1</sup> , Elmiro Vetere <sup>2</sup>	
- <i>Shape morphing using interpolation between implicit functions</i>	29
Daniil Ilyukhin	
- <i>The Fermat-Torricelli problem in normed spaces</i>	30
Denis Ilyutko	
- <i>Weight systems of framed chord diagrams: the Circuit-Nullity formula and Lie algebras</i>	31
Stefan Ivanov	
- <i>The Riemannian curvature identities on Almost Complex Calabi-Yau with torsion 6-manifold and generalized Ricci solitons</i>	32
Ismoilov Sherzod Shakirzhan ugli <sup>1</sup> , Artykbayev Artykbayev <sup>2</sup> , Jozef Mikeš <sup>3</sup>	
- <i>Geodesic mappings of semi-hyperbolic spaces</i>	34
Božidar Jovanović <sup>1</sup> , Tijana Šukilović <sup>2</sup> , Srdjan Vukmirović <sup>3</sup>	
- <i>Polynomial integrability of sub-Riemannian geodesic flows on compact Lie groups</i>	36
Ivan Limonchenko	
- <i>Persistent homology theory and toric topology</i>	37
Louis Kauffman	
- <i>Knot Dynamics and Vortex Reconnection</i>	39
Djordje Kocić	
- <i>The shape operator of real hypersurfaces in <math>S^6(1)</math></i>	40
Elena Kudryavtseva	
- <i>Generic bifurcations of tori in 3D integrable Hamiltonian systems</i>	41
Nenad Dj. Lazarov	
- <i>The velocity of one dimension cosmos</i>	43
Qing Li	
- <i>A continuum space is the infinitely great</i>	44
Katarina Lukić	
- <i>The Jacobi-orthogonality and Osserman tensors</i>	45
Elena N. Makhrova	
- <i>The structure of dendrites and dynamics of continuous maps on them</i>	46
Miroslav Maksimović <sup>1</sup> and Milan Zlatanović <sup>2</sup>	
- <i>Some curvature properties of quarter-symmetric metric connection</i>	48

Mancho Manev	
- <i>Ricci-Bourguignon Almost Solitons with Special Potential on Sasaki-like Almost Contact Complex Riemannian Manifolds</i> . . . . .	49
Miodrag Mateljević	
- <i>Lorentz Transformation and time dilatation</i> . . . . .	50
Jelena Matović <sup>1</sup> , Aleksandar Lipkovski <sup>2</sup>	
- <i>Calculation of quiver representations of finite ring digraphs</i> . . . . .	52
Josef Mikes <sup>1</sup> , Nadezda Guseva <sup>2</sup> , Irena Hinterleitner <sup>3</sup> , Patrik Peška <sup>4</sup> , Lenka Vítková <sup>5</sup>	
- <i>Geodesic mappings and their generalizations</i> . . . . .	53
Vladislava Milenković	
- <i>New types of mappings of generalized Riemannian spaces</i> . . . . .	56
Dmitry Millionshchikov	
- <i>Minimal models of nilmanifolds and complex structures</i> . . . . .	57
Velichka Milousheva	
- <i>Timelike Surfaces with Parallel Normalized Mean Curvature Vector Field and their Canonical Parameters</i> . . . . .	60
Svetislav M. Minčić	
- <i>On curvature tensors obtained by two non-symmetric linear connections</i> . . . . .	61
Alexandr Mishchenko	
- <i>Maslov index and infinitesimal Lagrangian manifolds</i> . . . . .	62
Mansi Mishra	
- <i>Weyl transform of a measure</i> . . . . .	65
Marija Najdanović	
- <i>On deformations preserving dual arc length in dual 3-space</i> . . . . .	66
Crina-Daniela Neacșu <sup>1</sup> , Gabriel-Eduard Vilcu <sup>2</sup>	
- <i>On the stability of the identity map of compact space forms</i> . . . . .	67
Yurii Nikonorov	
- <i>On geodesic orbit nilmanifolds</i> . . . . .	68
Yuri Nikolayevsky	
- <i>Killing tensors on symmetric spaces</i> . . . . .	69
Gleb Nosovski	
- <i>Implementation of Postprocessing Procedure of a Rapid Algorithm of Geometric Coding of Digital Images Using CUDA Architecture</i> . . . . .	70
Mustafa Özkan <sup>1</sup> and İrem Küpeli Erken <sup>2</sup>	
- <i>Fischer-Marsden Equation on Paracontact Geometry</i> . . . . .	71
Dhriti Sundar Patra	
- <i>Characterizations of Ricci-Bourguignon almost solitons</i> . . . . .	72
Milan Pavlović, Tijana Šukilović	
- <i>Integrability of the sub-Riemannian geodesic flow of the left-invariant metric on the Heisenberg group</i> . . . . .	73
Alexander Petkov	
- <i>Li-Yau sub-gradient estimates and Perelman-type entropy formulas for the heat equation in quaternionic contact geometry</i> . . . . .	74
Miloš Petrović	
- <i>Composition of conformal and projective mappings of generalized Riemannian spaces in Eisenhart's sense preserving certain tensors</i> . . . . .	75
Ariana Pitea	
- <i>Numerical algorithms on Hadamard manifolds</i> . . . . .	77

Theodore Popelensky	
- <i>On certain new results on the Steenrod algebra mod <math>p</math></i>	78
Ljiljana Radović	
- <i>The geometrical personalization of human organs 3D models by using the Characteristic Product Features methodology</i>	79
Lidija Rančić <sup>1</sup> and Svetozar Rančić <sup>2</sup>	
- <i>Basins of attractions of a new iterative method for finding simple zeros</i>	80
Pooja Rani and M.K. Vemuri	
- <i>The Brylinski beta function of a double layer</i>	81
Vladimir Rovenski	
- <i>Weak nearly Sasakian and weak nearly cosymplectic manifolds</i>	82
Vladimir Rovenski	
- <i>Willmore-type variational problem for foliated hypersurfaces</i>	84
Anton Savin	
- <i>On the Index Problem On Manifolds with Group Actions. Contributions of Fixed Points</i>	86
Anastasiiia Shubert	
- <i>Geometric Properties of the Inertia Tensor of a Rigid Body</i>	88
Dušan Simjanović <sup>1</sup> , Nenad Vesić <sup>2</sup>	
- <i>Invariants for geometric mappings</i>	89
Mića S. Stanković	
- <i>On some generalizations of Kahlerian spaces</i>	90
Sergey E. Stepanov	
- <i>New vanishing theorem of conformal Killing forms</i>	91
Jelena Stojanov	
- <i>Geometrical eigenproblem of various types higher order tensors in Riemannian space</i>	92
Milica Stojanović	
- <i>3-Triangulation of polyhedra and their connection graphs</i>	93
Alexey Tuzhilin	
- <i>Gromov-Hausdorff Distance and Geometric Optimization Problems</i>	94
Igor Uljarević	
- <i>Size in contact geometry</i>	95
Craig van Coevering	
- <i>Extremal Kähler metrics and the moment map</i>	96
Ana Velimirović <sup>1</sup> , Milan Zlatanović <sup>2</sup> ,	
- <i>Conformal transformations of curvature pseudotensors</i>	97
Ljubica Velimirović	
- <i>Curvature based shape consideration</i>	98
Nikola Velimirović <sup>1</sup> , Marija Najdanović <sup>2</sup> , Milan Zlatanović <sup>3</sup> , Andrija Zorić <sup>4</sup>	
- <i>Structural Shape Analysis</i>	99
Murali Vemuri	
- <i>A generalization of a result of Minakshisundaram and Pleijel</i>	100
Nenad Vesić <sup>1</sup> , Dušan Simjanović <sup>2</sup>	
- <i>New invariants for conformal mappings of Riemannian spaces</i>	101
Andrei Vesnin	
- <i>On Yamada polynomial of spatial graphs and Jones polynomial of related links</i>	102



Anton Vikhrov	
- <i>Geodesics in Gromov-Hausdorff class</i> . . . . .	104
Paweł Walczak	
- <i>Einstein Foliations</i> . . . . .	106
Kaori Yamaguchi	
- <i>On statistics which are almost sufficient from the viewpoint of the Fisher metrics</i> . . . . .	107
Handan Yıldırım	
- <i>Legendrian dual surfaces, lying in the 3-dimensional de Sitter space, of a spacelike curve in the 3-dimensional lightcone</i> . . . . .	108
Ekaterina Zhikhareva	
- <i>Semisimple algebraic Nijenhuis operators on small dimensional Lie algebras</i>	110
Filip Živanović	
- <i>(Non) existence of Lagrangians in hyperkähler manifolds</i> . . . . .	112



## PREFACE

The international conference XXII Geometrical Seminar 2024, is to be held at Vrnjačka Banja, Serbia from May 26th to 31st, 2024.

This Book of abstracts is collection of abstracts of talks to be presented at the conference from different geometric topics. Participants will have 15 and 30 minutes talks. The XXII Geometrical Seminar has more than 100 participants from all over the world. This meeting is bringing together mathematicians, physicists and engineers interested in Geometry and its applications. The aim of the Geometrical Seminar is to enable researches to give lectures on new results, exchange ideas, problems and conjectures. This issue contains abstracts of talks to be presented at the XXII Geometrical Seminar. The abstracts were accepted for presentation after having been subjected to the usual strict reviewing process of the conference committee. The editors thank the members of the Committees of Geometrical Seminar and to the Faculty of Sciences and Mathematics, Niš, Faculty of Mathematics, Belgrade, to the Faculty of Science, Kragujevac and Mathematical Institute SASA for their great effort in organizing this conference.

Editor

**Mića Stanković**

Department of Mathematics

Faculty of Science and Mathematics

University of Niš, Serbia

Email addres: `mica.stankovic@pmf.edu.rs`



## Integrable geodesic flows with rational first integrals

Sergei Agapov

Sobolev Institute of Mathematics SB RAS, Novosibirsk, Russia

E-mail: agapov.sergey.v@gmail.com

<https://orcid.org/0009-0009-4135-5792>

We study integrable geodesic flows on 2-surfaces. In most known examples, the additional integrals are polynomial in momenta. Polynomial integrals of small degrees are well- studied and classified in both local and global aspect of the problem.

On the other hand, as proved in [1], there exist local Riemannian metrics on 2-surfaces with integrable geodesic flows such that additional integrals are rational in momenta with any given degrees of a numerator and a denominator. However, constructing such examples in an explicit form turned out to be a very difficult problem. In this talk we will describe various methods and approaches which allowed us to construct rich families of 2-dimensional Riemannian metrics with integrable geodesic flows admitting rational first integrals (see [2], [3]).

The talk is based on joint work with V. Shubin.

### References

- [1] V.V. Kozlov, *On rational integrals of geodesic flows*, Regul. Chaotic Dyn., **19**, 601–606 (2014).
- [2] S. Agapov and V. Shubin, *Rational integrals of 2-dimensional geodesic flows: New examples*, Journ. Geom. Phys., **170**, 104389 (2021).
- [3] S. Agapov and V. Shubin, *New examples of non-polynomial integrals of two- dimensional geodesic flows*, J. Phys. A: Math. Theor., **57**, 015204 (2024).

## **Hopf Bundle, Listing's Law and Saccades**

**Dmitri Alekseevsky**

Institute for Information Transmission Problems RAS, Moscow, Russia

E-mail: dalekseevsky@iitp.ru

<https://orcid.org/0000-0002-6622-7975>

Our eyes continually move. Even while we fix our gaze on a still object, they participate in fixational eye movements – tremor, drift and microsaccades. The classical experiments by A. Yarbus show that compensation of eye movements leads to loss of vision in 2-3 sec.

The configuration space of the eye is described by the Listing's law. We propose an interpretation of the Listing's law in terms of the Hopf bundle as a Listing's hemisphere and use it for description of the saccades and saccadic cycles. We discuss the role of saccades and drift in vision. Roughly speaking, vision is the result of the interaction of information about light entering the retina and recorded by photoreceptors and oculomotor information about the eye position after remapping that accompanies a saccade.



## Neutrosophic Touchard Polynomials for Subclass of Analytic Functions

Khalifa Al Shaqsi

University of Nizwa, Oman  
E-mail: khalifa.alshaqsi@unizwa.edu.om  
<https://orcid.org/0009-0003-1184-9499>

By using the generalization of the neutrosophic Touchard Polynomials, we introduce a new subclass of analytic functions defined in the open unit disk. We then apply the  $q$ -Touchard Polynomials to investigate the estimates for the Taylor coefficients and Fekete-Szeg type inequalities of the functions belonging to this new subclass. In addition, we consider several corollaries and the consequences of the results presented in this paper. The neutrosophic Touchard Polynomials is expected to be significant in a number of areas of mathematics, science, and technology.

## **Convergence theorems for sequences of operator valued functions**

**Miloš Arsenović**

University of Belgrade, Faculty of Mathematics, Belgrade, Serbia

E-mail: [milos.arsenovic@matf.bg.ac.rs](mailto:milos.arsenovic@matf.bg.ac.rs)

<https://orcid.org/0000-0002-5450-2407>

We consider spaces of integrable operator valued functions, in the sense of Gelfand. Analogues of Dominated Convergence Theorem and of Vital Convergence Theorem are obtained, and a converse of Vitali Convergence Theorem is proved. Some of the results are specific for Hilbert space operators, namely those which rely on the notion of absolute value of operator. We note that earlier research centered on a single integrable operator valued functions and Radon - Nikodym properties. This is a joint work with Mihailo Krstic.

## On a coarse geometric approach to operator algebras

Andronick Arutyunov

Moscow Institute of Physics and Technologies, Russian Federation

E-mail: andronick.arutyunov@gmail.com

<https://orcid.org/0000-0002-6878-0993>

Coarse geometry is an interesting part of metric geometry, working with mappings that are not necessarily continuous and only need to be checked. In other words, there are such positive constants  $A, B$  that hold for metric spaces (e.g. graphs).

$$\forall x, y \in M_1 : \frac{1}{A} \rho^1(x, y) - B \leq \rho^2(f(x), f(y)) \leq A \rho^1(x, y) + B;$$

Here  $f : M_1 \rightarrow M_2$  is a mapping between metric spaces  $M_1, M_2$ . We will show that for graphs generated by the action of a group on itself (a well-known example are Cayley graphs), operator algebras are generated. For Cayley graphs this is the set of Fox derivations and for conjugacy diagrams we get a classical "Leibniz" derivation.

It is shown that the family of operators is coarsely invariant.

## Pointwise rectifying submanifolds and anti-torqued vector fields

Muhittin Evren Aydin

Firat University, Turkey  
E-mail: meaydin@firat.edu.tr

In this talk, we introduce a notion of pointwise rectifying submanifold in a Riemannian manifold with respect to a vector field called axis. We concern with the case that the axis is an anti-torqued vector field. Hence, we first find a necessary and sufficient condition for a Riemannian manifold to be endowed with an anti-torqued vector field. We then classify pointwise rectifying submanifolds when the axis is anti-torqued. Non-trivial examples are also provided.

This is a joint work with Adela Mihai and Cihan Özgür.

### References

- [1] B.-Y. Chen, *Differential geometry of rectifying submanifolds*, Int. Electron. J. Geom. 9(2) (2016), 18.
- [2] B.-Y. Chen, *Rectifying submanifolds of Riemannian manifolds and torqued vector fields*, Kragujev. J. Math. 41 (2017), 93103.
- [3] S. Deshmukh, I. Al-Dayel, D. M. Naik, *On an anti-torqued vector field on Riemannian*, Mathematics, 9 (2021), 2201.

## Left-invariant $f$ -structures on low-dimensional solvable Lie groups

Vitaly Balashchenko

Belarusian State University, Minsk, Belarus  
E-mail: balashchenko@bsu.by; vitbal@tut.by  
<https://orcid.org/0000-0002-9366-2411>

Metric  $f$ -structures by K.Yano ( $f^3 + f = 0$ ) play a leading role in the generalized Hermitian geometry (V.F.Kirichenko, D.Blair, S.Salamon and others). The most interesting classes are Killing  $f$ -structures ( $\nabla_X(f)(X) = 0$ ), nearly Kähler  $f$ -structures ( $\nabla_{fX}(f)(fX) = 0$ ), generalized nearly Kähler  $f$ -structures ( $f\nabla_{fX}(f)(fX) = 0$ ) introduced since the 1990s by V.F.Kirichenko, A.Gritsans, V.Balashchenko. In the particular case  $f = J$  ( $J^2 = -1$ ) all of them give the well-known nearly Kähler structures ( $\nabla_X(J)(X) = 0$ ). We also mention the class of Hermitian metric  $f$ -structures ( $T(X, Y) = 0$ ), where  $T(X, Y) = \frac{1}{4}f(\nabla_{fX}(f)(fY) - \nabla_{f^2X}(f)(f^2Y))$  is the composition tensor (V.F.Kirichenko).

We concentrate on left-invariant metric  $f$ -structures on solvable Lie groups in dimension 3 and 4. Note that three-dimensional solvable Lie algebras were first classified by G.M.Mubarakzyanov in 1963. Using special left-invariant metric  $f$ -structures on all three-dimensional solvable Lie algebras we present the examples of all the classes in the generalized Hermitian geometry above mentioned. In particular, for the Lie algebras  $[e_1, e_2] = \lambda e_2 - e_3$ ;  $[e_1, e_3] = e_2 + \lambda e_3$  depending on parameter  $\lambda$  we define  $f$ -structure by the requirement  $f(e_1) = 0, f(e_2) = -e_3, f(e_3) = e_2$ . It was proved that the left-invariant  $f$ -structure on the corresponding Lie group is a nearly Kähler  $f$ -structure for  $\lambda = -1$  and a Hermitian  $f$ -structure for arbitrary  $\lambda$ . Among four-dimensional solvable Lie groups the oscillator group is of especial interest in geometry, control theory and other applications.

The corresponding Lie algebra is defined by the commutator relations  $[e_2, e_3] = e_1$ ,  $[e_2, e_4] = -e_3$ ,  $[e_3, e_4] = e_2$ . We completely investigate all basic left-invariant  $f$ -structures of rank 2 on this Lie group in the above sense.

This is a joint work with Victoria Kunitsa.



## The mod $p$ Buchstaber invariant

Djordje Baralić

Mathematical Institute SANU, Belgrade, Serbia

E-mail: [djbaralic@mi.sanu.ac.rs](mailto:djbaralic@mi.sanu.ac.rs)

<https://orcid.org/0000-0003-2836-7958>

In this talk, we present combinatorial and topological properties of the universal complexes  $X(\mathbb{F}_p^n)$  and  $K(\mathbb{F}_p^n)$  whose simplices are certain unimodular subsets of  $\mathbb{F}_p^n$ . We calculate their  $\mathbf{f}$ -vectors and the bigraded Betti numbers of their Tor-algebras, show that they are shellable, and find their applications in toric topology and number theory. We showed that the Lusternick-Schnirelmann category of the moment angle complex of  $X(\mathbb{F}_p^n)$  is  $n$ , provided  $p$  is an odd prime and the Lusternick-Schnirelmann category of the moment angle complex of  $K(\mathbb{F}_p^n)$  is  $\lfloor \frac{n}{2} \rfloor$ . Based on the universal complexes, we introduce the Buchstaber invariant  $s_p$  for a prime number  $p$ . We investigate the mod  $p$  Buchstaber invariant of the skeleta of simplices for a prime number  $p$  and compare them for different values of  $p$ . For  $p = 2$ , the invariant is the real Buchstaber invariant. Our findings reveal that these values are generally distinct. Additionally, we determine or estimate the mod  $p$  Buchstaber invariants of some universal simplicial complexes  $X(\mathbb{F}_p^n)$ . The talk is based on joint research with Aleš Vavpetić and Aleksandar Vučić.

## Almost complex manifolds with Norden metrics

**Cornelia-Livia Bejan**

"Gh. Asachi" Technical University of Iasi, Romania

E-mail: bejanliv@yahoo.com

<https://orcid.org/0000-0001-6963-7710>

On an almost complex manifold, a Norden structure is a semi- Riemannian metric, with respect to which the almost complex structure is an anti-isometry. We construct here a class of almost complex manifolds with Norden metric on the total space of the cotangent bundle (of a manifold with a symmetric connection), endowed with a natural Riemann extension (which generalize the Riemann extension, introduced by Patterson, Walker). Then we give necessary and sufficient conditions to belong to certain classes of Ganchev-Borisov classification of almost complex manifolds with Norden metric. Some harmonicity problems are studied. At the end, we construct a class of hypercomplex Norden structures.

## Remarks on weighted Einstein manifolds

Miguel Brozos-Vázquez

CITMAga-Universidade da Coruña, A Coruña, Spain

E-mail: miguel.brozos.vazquez@udc.es

<https://orcid.org/0000-0003-4945-9587>

A Riemannian manifold  $(M, g)$  is endowed with a smooth density function  $f$ , and two real parameters  $m$  and  $\mu$ , giving rise to a smooth metric measure space  $(M, g, f, m, \mu)$ . In order to analyze smooth metric measure spaces, it is useful to consider weighted objects such as the weighted Schouten tensor  $P_f^m$  or the weighted Weyl tensor  $W_f^m$ , which reflect different aspects of the influence of the density on the geometry of the underlying manifold.

Weighted Einstein manifolds are smooth metric measure spaces which satisfy  $P_f^m = \lambda g$ , for  $\lambda$  constant. This is a generalization of Riemannian Einstein manifolds. However, although usual Einstein metrics have harmonic Weyl tensor, this is no longer true in the weighted context. Motivated by this fact, we analyze which smooth metric measure spaces are weighted Einstein and have harmonic Weyl tensor (in a suitable weighted sense). First, we provide a local classification result showing that, under the harmonicity condition, the underlying manifold is Einstein or decomposes as a particular warped product. Moreover, if the manifold is complete, then we provide a global rigidity theorem which shows that it either is a weighted analogue of a space form, or it belongs to a particular family of Einstein warped products with  $\lambda < 0$ .

## Great antipodal sets and applications

**Bang-Yen Chen**

Michigan State University, United States

E-mail: [chenb@msu.edu](mailto:chenb@msu.edu)

<https://orcid.org/0000-0002-1270-094X>

Greatest antipodal sets were introduced in 1982 by Chen and Nagano. Since then some applications of greatest antipodal sets to several areas of mathematics have been established by various mathematicians. In this talk, I will discuss some recent applications of greatest antipodal sets.

## The Willmore energy variations and energy efficient architecture

Milica Cvetković<sup>1</sup>, Ana Mitrović<sup>2</sup>

<sup>1</sup>The Academy of Applied Technical and Preschool Studies, Niš, Serbia

E-mail: milica.cvetkovic@akademijanis.edu.rs

<https://orcid.org/0000-0003-0418-2000>

<sup>2</sup>University of Belgrade, Faculty of Architecture, Belgrade, Serbia

[ana@ateljedv.com](mailto:ana@ateljedv.com)

Since the shape is an important feature of objects and can be immensely useful in characterizing objects, it should point out the shape analysis considering the variation of magnitudes that characterize the shape itself. One of the fundamental functionals that measure the bending of a surface is the Willmore energy. In view of the meaning of ruled surfaces in aesthetics, statics, scale and manufacturing technologies, we point out the possibility of a mathematical analysis in the case of infinitesimal deformations by considering the variations of the Willmore energy on Gaudi surface under infinitesimal bending in  $R^3$ . Application could be connected with energy efficient building design.

**AMS Subj. Class.:** 53A05, 53C45, 92C40, 51M15

**Key words:** Willmore energy, Gaudi surface, Infinitesimal bending, Variation, Energy efficient architecture.

## Almost Kähler structures on complex hyperbolic space

Andrijana Dekić

Mathematical Institute of Serbian Academy of Sciences and Arts, Belgrade, Serbia

E-mail: andrijanadekic@turing.mi.sanu.ac.rs

<https://orcid.org/0000-0001-6955-0930>

In this talk we consider Complex hyperbolic space as a solvable Lie group. Based on the recent classification of Riemannian left-invariant metrics [1] we describe all left-invariant almost Kähler structures on that group [2]. We calculate the canonical connection and obtain the characterization by its torsion tensor.

### References

- [1] A. Dekić, M. Babić, S. Vukmirović, *Classification of Left Invariant Riemannian metrics on Complex hyperbolic space*, Mediterr. J. Math., 19:232, (2022)
- [2] A. Dekić, *Almost Kähler structures on complex hyperbolic space*, Filomat, Vol 37, No 25, (2023)



## Curvature properties of pseudosymmetry type of some 2-quasi Einstein manifolds

Ryszard Deszcz<sup>1</sup>, Małgorzata Głogowska<sup>2</sup>, Jan Jełowicki<sup>3</sup>,  
Miroslava Petrović-Torgašev<sup>4</sup>, and Georges Zafindratafa<sup>5</sup>

<sup>1</sup> Department of Applied Mathematics, Wrocław University of Environmental and Life Sciences, Wrocław, Poland

E-mail: Ryszard.Deszcz@upwr.edu.pl

<https://orcid.org/0000-0002-5133-5455>

<sup>2</sup> Department of Applied Mathematics, Wrocław University of Environmental and Life Sciences, Wrocław, Poland

E-mail: Malgorzata.Glogowska@upwr.edu.pl

<sup>3</sup> Department of Applied Mathematics, Wrocław University of Environmental and Life Sciences, Wrocław, Poland

E-mail: Jan.Jelowicki@upwr.edu.pl

<sup>4</sup> Department of Sciences and Mathematics, State University of Novi Pazar, Serbia

E-mail: mirapt@kg.ac.rs

<sup>5</sup> Laboratoire de Mathématiques pour l'Ingénieur (LMI), Université Polytechnique Hauts-de-France,

E-mail: Georges.Zafindratafa@uphf.fr

In this talk we present curvature properties of pseudosymmetry type of 2-quasi Einstein semi-Riemannian manifolds  $(M, g)$ ,  $\dim M \geq 4$ , with the Riemann-Christoffel curvature tensor  $R$  expressed by a linear combination of Kulkarni-Nomizu products  $g \wedge g$ ,  $g \wedge S$ ,  $S \wedge S$  and  $g \wedge S^2$ , formed by the metric tensor  $g$ , the Ricci tensor  $S$  and its square  $S^2$ . Our talk is based on [1], [2], [3] and [4].

### References

- [1] R. Deszcz, M. Głogowska, M. Hotłoś, M. Petrović-Torgašev and G. Zafindratafa, *A note on some generalized curvature tensor*, Int. Electron. J. Geom. 16 (1) (2023), 379–397.
- [2] R. Deszcz, M. Głogowska, M. Hotłoś, and K. Sawicz, *Hypersurfaces in space forms satisfying a particular Roter type equation*, arXiv: 2211.06700v3 [math.DG] 22 Nov 2023, 24 pp.

- [3] R. Deszcz, M. Głogowska, J. Jełowicki, M. Petrović-Torgašev, G. Zafindratafa, *Curvature properties of pseudosymmetry type of some 2-quasi Einstein manifolds*, to appear.
- [4] R. Deszcz, M. Głogowska, J. Jełowicki, G. Zafindratafa, *Curvature properties of some class of warped product manifolds*, Int. J. Geom. Meth. Modern Phys. 13 (2016), 1550135 (36 pages).

## The Schwarzschild-de Sitter Metric of Nonlocal $\sqrt{dS}$ Gravity

Ivan Dimitrijević

University of Belgrade, Faculty of Mathematics, Belgrade, Serbia

E-mail: [ivan.dimitrijevic@matf.bg.ac.rs](mailto:ivan.dimitrijevic@matf.bg.ac.rs)

<https://orcid.org/0000-0003-3092-2295>

Nonlocal de Sitter gravity model defined by the action

$$S = \frac{1}{16\pi G} \int (R - 2\Lambda + P(R) \mathcal{F}(\square) Q(R)) \sqrt{-g} dx^4, \quad (1)$$

which we denote as  $\sqrt{dS}$  gravity, contains an exact vacuum cosmological solution which mimics dark energy and dark matter and is in very good agreement with the standard model of cosmology. In this presentation we investigate how it works at galactic scale. We consider Schwarzschild-de Sitter metric of the  $\sqrt{dS}$  gravity model. We present an approximative solution of linearized equation, which is related to space metric far from the massive body, where gravitational field is weak. The obtained solution is of particular interest for examining the possible role of non-local de Sitter gravity  $\sqrt{dS}$  in describing the effects in galactic dynamics that are usually attributed to dark matter. The solution has been tested on the Milky Way and the spiral galaxy M33 and is in good agreement with observational measurements.

## **Yamabe solitons on submanifolds of Kähler manifolds**

**Miloš Djorić**

University of Belgrade, Faculty of Mathematics, Belgrade, Serbia

E-mail: milos.djoric@matf.bg.ac.rs

<https://orcid.org/0000-0001-5462-5522>

Yamabe solitons, defined by R. Hamilton, are the self-similar solutions to the Yamabe flow, which evolves an initial metric on given Riemannian manifold to a new one with constant scalar curvature, within the same conformal class of metrics. For any such soliton, there exists the soliton vector field which relates the Lie derivative of the metric with the scalar curvature.

We study submanifolds of Kähler manifolds which admit (quasi-)Yamabe soliton structure, especially CR submanifolds with the Reeb vector field as a soliton vector field. When the ambient manifold is a non-flat complex space form, we give a complete classification of such solitons, under certain conditions.

This is a joint work with Mirjana Djorić.

## Nonlocal de Sitter Gravity $\sqrt{dS}$ and the Dark Side of the Universe

Branko Dragovich

Institute of Physics, University of Belgrade, Belgrade, Serbia  
Mathematical Institute, Serbian Academy of Sciences and Arts, Belgrade, Serbia  
E-mail: dragovich@ipb.ac.rs

<https://orcid.org/0000-0002-5818-0150>

I will present nonlocal de Sitter gravity model, denoted by  $\sqrt{dS}$ ,

$$S = \frac{1}{16\pi G} \int d^4x \sqrt{-g} (R - 2\Lambda + \sqrt{R - 2\Lambda} \mathcal{F}(\square) \sqrt{R - 2\Lambda}), \quad (2)$$

where  $\Lambda$  is the cosmological constant,  $\square$  is d'Alembert-Beltrami operator, and nonlocal operator  $\mathcal{F}(\square)$  has the following general form:

$$\mathcal{F}(\square) = \sum_{n=1}^{+\infty} (f_n \square^n + f_{-n} \square^{-n}). \quad (3)$$

This nonlocal model is unique compared to other non-local models and its properties will be pointed out.

Several exact vacuum cosmological solutions are obtained. One of these solutions is  $a(t) = At^{\frac{2}{3}} e^{\frac{\Lambda}{14} t^2}$ , and it mimics effects that are usually assigned to dark matter and dark energy. Obtained results are in a very good agreement with the standard model of cosmology. Some other solutions are examples of the nonsingular bounce ones in flat, closed and open universe. There are also singular and cyclic solutions. All these cosmological solutions are a result of nonlocality and do not exist in the local de Sitter case.

This talk is based on the joint research with Ivan Dimitrijevic, Jelena Stankovic and Zoran Rakic, and our common paper: JHEP 12 (2022) 054.

## How to connect two most beautiful geometric theorems?

Vladimir Dragović

University of Texas at Dallas, MISANU

E-mail: vladad@turing.mi.sanu.ac.rs

<https://orcid.org/0000-0002-0295-4743>

Using Marden's Theorem from geometric theory of polynomials, we show that for every triangle there is a unique ellipse such that the triangle is a billiard trajectory within that ellipse. Since 3-periodic trajectories of billiards within ellipses are examples of the Poncelet polygons, our considerations provide a new insight into the relationship between Marden's Theorem and the Poncelet Porism, two gems of exceptional classical beauty. We also show that every parallelogram is a billiard trajectory within a unique ellipse. We prove a similar result for the self-intersecting polygonal lines consisting of two pairs of congruent sides, named "Darboux butterflies". In each of three considered cases, we effectively calculate the foci of the boundary ellipses. This is based on joint work with Milena Radnovic.

### References

- [1] V. Dragović, M. Radnović, *Is every triangle a trajectory of an elliptical billiard?* arXiv 2405.08922, 2024
- [2] V. Dragović, *Poncelet-Darboux curves, their complete decomposition and Marden theorem*, International Math. Res. Notes, 2011, Vol. 2011, p. 3502-3523.

## **Finsler geodesic orbit metrics**

**Zdenek Dusek**

Institute of Technology and Business in Ceske Budejovice, Czech Republic  
E-mail: [zdusek@mail.vstecb.cz](mailto:zdusek@mail.vstecb.cz)

There are many results about geodesic orbit metrics in Riemannian geometry. Recently, the topic attracted attention also in Finsler geometry. Our approach to geodesic orbit metrics is through geodesic graphs, which are the generating sets for homogeneous geodesics. In Riemannian geometry the interesting cases are those which have nonlinear structure. Various special Finsler metrics arising from Riemannian geodesic orbit metrics modify the structure of geodesic graphs in particular way. We are able to prove that some of these Finsler metrics have also geodesic orbit property, for example the  $(\alpha, \beta)$  metrics. The examples of Finslerian geodesic graphs on nilmanifolds and on spheres will be presented. Part of the talk is based on the joint work with Teresa Arias-Marco.

## On maps obtained by small perturbations of skew products

Lyudmila Efremova

Nizhny Novgorod State University, Nizhny Novgorod, Russia

E-mail: [lefunn@gmail.com](mailto:lefunn@gmail.com)

<https://orcid.org/0000-0001-5821-6697>

We consider  $C^1$ -smooth maps obtained by small perturbations (in the  $C^1$ -norm) of  $C^1$ -smooth skew products on a compact cylinder. We prove the geometric integrability of these maps and construct the example of the family of such maps with the global attractor, which is one-dimensional ramified continuum with the set of the continuum cardinality of ramification points of the order 3.

This research is supported by Russian Science Foundation (RSF), grant 24-21-00242, <https://rscf.ru/en/project/24-21-00242/>.



## Some Remarks on Quasi-para-Sasakian Manifolds

İrem Küpeli Erken

Bursa Technical University, Faculty of Engineering and Natural Science,  
Department of Mathematics, Bursa, Turkey  
E-mail: [irem.erken@btu.edu.tr](mailto:irem.erken@btu.edu.tr)  
<https://orcid.org/0000-0003-4471-3291>

In this talk, we give structure and curvature identities for quasi-para-Sasakian manifolds. We investigate D-homothetic deformations on quasi-para-Sasakian manifolds and we characterize them according to constant curvature. Finally, we give a method for constructing examples of three-dimensional conformally flat quasi-para-Sasakian manifolds.

## **Spectrum problem of Schrodinger operators related to Landau-Ginzburg models**

**FAN Huijun**

School of Mathematical Sciences, Peking University, China  
E-mail: fanhj@math.pku.edu.cn

An LG model  $(M, f)$  is given by a noncompact complex manifold  $M$  and the holomorphic function  $f$  defined on it, which is an important model in string theory. Recently, LG models attracted more attention due to their important applications in mirror symmetry and complex geometry. Typical examples include Laurent polynomials defined on log Calabi-Yau manifolds, which are mirror to different Fano manifolds. The B-theory of a LG model is related to the Hodge theory of a twisted Laplacian operator, which is a form Schrodinger operator defined on the noncompact manifold  $M$ . This talk will concern the spectrum problem of this kind of Schrodinger operators. We will prove a fundamental spectrum theorem when  $(M, f)$  satisfies a strongly tame condition. This theorem can be applied to many LG models appearing in mirror symmetry.

## **Critical metrics for quadratic curvature functionals**

**Eduardo Garcia-Rio**

University of Santiago de Compostela, Faculty of Mathematics, Santiago de Compostela, Spain

E-mail: [eduardo.garcia.rio@usc.es](mailto:eduardo.garcia.rio@usc.es)

<https://orcid.org/0000-0003-1195-1664>

The purpose of this talk is to describe homogeneous critical metrics for quadratic curvature functionals in dimensions three and four. In addition to Einstein metrics, there are a number of non-symmetric examples which, in some cases, correspond to Ricci solitons.

## **Applications of algebraic topology to operator algebras: Homogeneous $C^*$ -algebras**

**Ilja Gogić**

University of Zagreb Department of Mathematics, Faculty of Science, Bijenicka 30,  
10000 Zagreb, Croatia  
E-mail: [ilja@math.hr](mailto:ilja@math.hr)  
<https://orcid.org/0000-0003-4445-8163>

The class of commutative  $C^*$ -algebras was fully described in the early days of the development of the theory of operator algebras: these are precisely algebras of continuous complex-valued functions on locally compact Hausdorff spaces that vanish at infinity. After commutative, the next simplest class are the homogeneous  $C^*$ -algebras, i.e.  $C^*$ -algebras whose all irreducible representations are of the same finite dimension. However, the problem of classification of such algebras becomes considerably more complicated due to certain topological phenomena which do not occur in the commutative case. In this talk we shall present the overview of homogeneous  $C^*$ -algebras with an emphasis on some more recent results.

## Shape morphing using interpolation between implicit functions

Admir Huseini<sup>1</sup>, Elmiro Vetere<sup>2</sup>

<sup>1</sup>3Shape and Institute of Mathematics, University of St. Cyril and Methodius,  
Skopje, North Macedonia  
E-mail: huseini@risat.org

<sup>2</sup> 3Shape, Copenhagen, Denmark

In this paper, we present a deformation algorithm based on combinatorial or piecewise linear triangulated manifolds. Given two combinatorial manifolds  $M_1$  and  $M_2$ , that are not necessarily homeomorphic, we construct a deformation mapping that is approximately continuous and resolves the issues that arise due to the discontinuity in topology and geometry.

The technique we use is based on constructing an interpolation between level set representations of combinatorial manifolds. We describe in detail the process of generating a level set or implicit function from a combinatorial manifold and the inverse process of obtaining a combinatorial manifold from a level set.

We show that the interpolation between the level sets can be interpreted as the level set of a higher dimensional manifold  $M$ , which contains  $M_1$  and  $M_2$  as boundaries. It contains important geometric information about the discontinuities of the deformation. As it is known from Morse theory, they appear exactly at the critical points on  $M$ . Each intermediate manifold  $M_t$  corresponds to a cross-section of  $M$  with a hyperplane.

We give applications in computer graphics in case the manifolds have dimensions 1, 2 or 3 and discuss efficient ways how to implement it in the graphics processing unit or GPU.

## The Fermat-Torricelli problem in normed spaces

Daniil Ilyukhin

Lomonosov Moscow State University, Moscow, Russian Federation

E-mail: daniil.ilukhin@math.msu.ru

<https://orcid.org/0000-0001-7744-3991>

The talk discusses a generalization of the classical Fermat- -Torricelli problem to normed spaces of arbitrary finite dimension.

The problem of finding a point that minimizes the sum of distances from it to a given set of points in a metric space was first mentioned in the 17th century. In 1643 Fermat posed a problem for three points on the Euclidean plane, and in the same century Torricelli proposed a solution to this problem. Since then, various generalizations of this problem have been considered. The problem was formulated for an arbitrary number of points, the dimension of the space, as well as the norm given in this space. The simplicity of the formulation allows us to consider the problem even in an arbitrary metric space.

The work describes the application of a geometric approach to the problem and presents some new results that are obtained in the framework of real finite-dimensional normed spaces, called Minkowski spaces.

The aim of this talk is to present necessary and sufficient conditions for the uniqueness of the solution of the Fermat- -Torricelli problem for any  $n$  points in a fixed space, and more precise conditions for normed planes and three- dimensional spaces.

In addition, examples of the application of the criterion in the norms given by regular polygons and regular polyhedra are given.

## **Weight systems of framed chord diagrams: the Circuit-Nullity formula and Lie algebras**

**Denis Ilyutko**

Chair of Differential Geometry and Applications  
Faculty of Mechanics and Mathematics, Lomonosov Moscow State University  
Leninskie Gory 1, Moscow, 119991 Russia  
E-mail: [denis.ilyutko@math.msu.ru](mailto:denis.ilyutko@math.msu.ru)  
<https://orcid.org/0000-0003-3184-0494>

Chord diagrams and 4T-relations appear in studying Vassiliev invariants (finite-order invariants) of knots. The theory of Vassiliev invariants is tightly associated with the theory of finite-order J-invariants of planar curves that was developed by Arnold. In studying this theory, there arise the notions of framed chord diagram and framed 4T relation. Framed chord diagrams are also applied in describing the combinatorics of non-general position of Legendre knots in three-dimensional manifolds.

One of the main properties of chord diagrams is as follows: considering the formal linear combinations of chord diagrams modulo 4T-relations, we can rather simply define the product of these linear combination. As a result, we obtain the algebra of chord diagrams. According to the Vassiliev-Kontsevich theorem, weight systems, that is, linear functions on this algebra, lead to Vassiliev invariants of knots. However, in the case of framed chord diagrams, taking a connected sum is not a correct operation.

In view of this, there arises the problem of searching for invariants of framed chord diagrams that essentially distinguish chords of different framings. In our work we generalize the Bar-Natan construction of weight systems induced by Lie algebra representations to the case of framed chord diagrams. This is a joint work with I.M.Nikonov.

## **The Riemannian curvature identities on Almost Complex Calabi-Yau with torsion 6-manifold and generalized Ricci solitons**

**Stefan Ivanov**

Sofia University "St.Kl.Ohridski" and IMI-BAS, Sofia, Bulgaria  
E-mail: ivanovsp@fmi.uni-sofia.bg

Curvature properties of the torsion connection on an Almost complex CYT 6-manifold are investigated. It is observed that on a compact ACYT 6-manifold the Nijenhuis tensor is parallel with respect to the torsion connection. It is shown that a compact ACYT 6-manifold has closed torsion if and only if the Ricci tensor of the torsion connection is equal to the covariant derivative of the Lee form with respect to the torsion connection. It is proved that a compact ACYT 6-manifold with closed torsion is Ricci flat if and only if either the norm of the torsion is constant or the Riemannian scalar curvature is constant. It is observed that any compact ACYT 6-manifold with closed torsion 3-form is a generalized gradient Ricci soliton and this is equivalent to a certain vector field to be parallel with respect to the torsion connection. In particular, this vector field is an infinitesimal automorphism of the  $SU(3)$  structure.

It is also proved that on a compact ACYT 6-manifold the curvature of the torsion connection  $R \in S^2\Lambda^2$  and has vanishing Ricci tensor if and only if the three-form torsion is parallel with respect to the Levi-Civita and to the torsion connection simultaneously. In particular, the conditions  $R \in S^2\Lambda^2, Ric = 0$  are equivalent to the condition that the curvature of the torsion connection satisfies the Riemannian first Bianchi identity, i.e. it is Kähler-like. In this case the torsion 3-form is closed and co-closed.



In the complex case, it follows that the curvature of the Strominger-Bismut connection on a six dimensional CYT space satisfies the conditions  $R \in S^2\Lambda^2, Ric = 0$  if and only if it is Kähler-like, therefore flat, and has parallel torsion with respect to both the Levi-Civita and Strominger-Bismut connections.

The research is partially supported by Contract KP-06-H72-1/05.12.2023 with the National Science Fund of Bulgaria and the National Science Fund of Bulgaria, National Scientific Program "VIHREN", Project KP-06-DV-7.

## Geodesic mappings of semi-hyperbolic spaces

Ismoilov Sherzod Shakirzhan ugli<sup>1</sup>, Artykbayev Artykbayev<sup>2</sup>,  
Jozef Mikeš<sup>3</sup>

<sup>1</sup>Tashkent State Transport University, Uzbekistan  
E-mail: Sh.ismoilov@nuu.uz

<sup>2</sup>Tashkent State Transport University, Uzbekistan  
E-mail: aartykbaev@mail.ru

<sup>3</sup>Palacky University Olomouc, Czech Republic  
E-mail: josef.mikes@upol.cz  
<https://orcid.org/0000-0003-2098-7976>

In the monograph by Rosenfeld, definitions of multidimensional semi-hyperbolic spaces with projective metrics are given [1]. A. Artykbaev defines semi hyperbolic spaces as geometry on semi-Euclidean spaces [2]. We will construct a geodesic mapping [3] of semi-hyperbolic spaces to semi-Euclidean spaces  ${}^{l_1 l_2} R_n^m$ .

The following statements are true:

1. Let  $TX = \frac{X + (X, l_0)}{(X, l_0)}$  be the mapping  ${}^{l_1 l_2} S_n^m$  in  ${}^{l_1 l_2} R_n^m$ , where  $X \in {}^{l_1 l_2} R_n^m$ ,  $\{l_0, l_1, \dots, l_n\}$  is the basis of the space, and  $(X, l_0)$  is the scalar product. Then  $TX$  is a geodesic mapping.
2. The semi-hyperbolic space  ${}^{l_1 l_2} S_n^m$  is geodesically mapped to the inner sphere  ${}^{l_1 l_2} S_{n-1}^m \subset {}^{l_1 l_2} R_n^m$  of the semi-Euclidean space.
3. When mapping  $TX$ , the order of the metric is preserved.

The resulting mapping will be generalized to the Cayley-Klein interpretation for semi-hyperbolic spaces.

## References

- [1] B.A. Rosenfeld, *Non-Euclidean spaces*. Moscow: Nauka, 1969.
- [2] A. Artikbaev, D. Sokolov, *Geometry in general in flat space-time*. Tashkent: Fan, 1991.
- [3] J. Mikeš et al, *Differential geometry of special mappings*. Olomouc: Palacky Univ. Press, 2019.

## Polynomial integrability of sub-Riemannian geodesic flows on compact Lie groups

Božidar Jovanović<sup>1</sup>, Tijana Šukilović<sup>2</sup>, Srdjan Vukmirović<sup>3</sup>

<sup>1</sup>Mathematical Institute, Serbian Academy of Sciences and Arts, Kneza Mihaila  
36, 11000 Belgrade, Serbia

E-mail: bozaj@mi.sanu.ac.rs

<https://orcid.org/0000-0002-3393-4323>

<sup>2,3</sup>Faculty of Mathematics, University of Belgrade, Studentski trg 16, 11000  
Belgrade, Serbia

E-mail: tijana.sukilovic@matf.bg.ac.rs, srdjan.vukmirovic@matf.bg.ac.rs

<sup>2</sup> <https://orcid.org/0000-0001-6371-3081>

We classify almost multiplicity free subgroups  $K$  of compact simple Lie groups  $G$ . The problem is related to the integrability of Riemannian and sub-Riemannian geodesic flows of left-invariant metrics defined by a specific extension of integrable systems from  $T^*K$  to  $T^*G$ .

We use the chains of subalgebras in order to construct integrable non-holonomic and sub-Riemannian flows with left-invariant metrics and left-invariant nonholonomic distributions on compact Lie groups. While the nonholonomic problem is not Hamiltonian, the sub-Riemannian is.

The talk is based on the paper *Almost multiplicity free subgroups of compact Lie groups and polynomial integrability of sub-Riemannian geodesic flows* by Božidar Jovanović, Tijana Šukilović, Srdjan Vukmirović [Letters in Mathematical Physics, 114 (1) Article no. 14, 2024].

## Persistent homology theory and toric topology

Ivan Limonchenko

Mathematical Institute of the Serbian Academy of Sciences and Arts, Belgrade,  
Serbia

E-mail: ilimonchenko@gmail.com

<https://orcid.org/0000-0002-2072-8475>

Toric topology is a new rapidly developing area of Mathematics in the crossroads of algebraic geometry, homotopy theory, equivariant topology, homological algebra and combinatorics. The main object of study in toric topology is a polyhedral product.

Given any topological pair  $(X, A)$  and simplicial complex  $K$  on  $[m] := \{1, 2, \dots, m\}$ , the polyhedral product  $(X, A)^K$  is a subset in  $X^m$  s.t. an  $m$ -tuple  $(x_1, \dots, x_m)$  is in  $(X, A)^K$  iff there is a  $\sigma \in K$ :  $i \notin \sigma \Rightarrow x_i \in A$ . Our research is based on the use of the moment-angle complex  $\mathcal{Z}_K := (D^2, S^1)^K$  of  $K$ .

Various applications of toric topology are related with the theorem by Baskakov, Buchstaber and Panov stating that cohomology  $H^*(\mathcal{Z}_K; \mathbb{k})$  of  $\mathcal{Z}_K$  with coefficients in a PID with unit  $\mathbb{k}$  is isomorphic as a  $\mathbb{k}$ -algebra to the Tor-algebra  $\text{Tor}_{\mathbb{k}[m]}(\mathbb{k}[K], \mathbb{k})$  of  $K$  (homology of the Koszul complex of  $K$ ), where  $\mathbb{k}[m] := \mathbb{k}[v_1, \dots, v_m]$  and the Stanley-Reisner ring  $\mathbb{k}[K]$  is its quotient by the ideal generated by all monomials corresponding to the non-simplices in  $K$ .

In this talk we shall discuss certain key results obtained recently in the framework of the persistent double homology theory.

Namely, we shall introduce three equivalent definitions (geometrical, algebraic and topological ones) of a new differential acting on the singular (co)chain complex of  $\mathcal{Z}_K$  and define the double cohomology of  $\mathcal{Z}_K$  w.r.t. the arising bicomplex. Then we shall discuss the applications of this theory in topological data analysis, as well as open problems and conjectures.

The talk is based on the recent projects j.w. Anthony Bahri (Rider University, USA), Taras Panov (HSE University, Russia), Jongbaek Song (Pusan National University, Korea) and Donald Stanley (University of Regina, Canada).

## **Knot Dynamics and Vortex Reconnection**

**Louis Kauffman**

University of Illinois at Chicago, United States

E-mail: [loukau@gmail.com](mailto:loukau@gmail.com)

<https://orcid.org/0000-0003-4135-8685>

This talk will discuss how one dimensional vortices in a fluid reconnect to either increase or decrease their topological complexity. We will give examples via photography and computer graphics and we shall discuss the problem of finding the least number of reconnections that unknot a given configuration.

## The shape operator of real hypersurfaces in $S^6(1)$

Djordje Kocić

Faculty of Mathematics, University of Belgrade, Serbia

E-mail: [djordje.kocic@matf.bg.ac.rs](mailto:djordje.kocic@matf.bg.ac.rs)

<https://orcid.org/0000-0003-2255-2992>

We will present two results concerning real hypersurfaces in the six-dimensional sphere  $S^6(1)$ . More precisely, we prove that real hypersurfaces with Lie-parallel shape operator  $A$  must be totally geodesic hyperspheres. Additionally, we classify real hypersurfaces in nearly Kähler sphere  $S^6(1)$  whose Lie derivative of shape operator coincides with its covariant derivative.



## Generic bifurcations of tori in 3D integrable Hamiltonian systems

Elena Kudryavtseva

Moscow State University, Russia  
Moscow 119991, Leninskie gory, 1, MSU  
E-mail: ekudr@list.ru  
<https://orcid.org/0000-0002-3745-2663>

We study singularities of the Lagrangian fibration given by a completely integrable Hamiltonian system with 2 or 3 degrees of freedom. Our purpose is to study (degenerate) singularities of ‘typical’ integrable systems.

As a particular case, one can study bifurcations of equilibria in 1-parameter families of integrable systems with 2 degrees of freedom. In this way, Lerman and Umanskii (1981) introduced the so-called parabolic singularity (which is the simplest degenerate singularity of generic integrable systems) and proved that it is structurally stable under (small enough) integrable perturbations of the system.

Another approach is to consider special integrable Hamiltonian systems (called semi-toric), in which the additional first integral generates a Hamiltonian circle action. By combining these two approaches, Duistermaat (1984) and van der Meer (1985) studied the Hamiltonian Hopf bifurcation and its analogue with resonances (that are the simplest generic bifurcations of degenerate equilibria). They proved that such bifurcations appear in generic 1-parameter families of semi-toric integrable systems with 2 degrees of freedom.

A third approach is to study compact periodic orbits instead of bifurcations of relative equilibria (thus the multiplicities of the periodic orbit can be different from 1, allowing a ‘twist’ of the orbit). By combining this approach with the previous one (concerning the existence of a circle action), Kalashnikov (1998) described an infinite series of singularities called parabolic orbits with resonances, and proved that they appear in generic 1-parameter families of semi-toric integrable systems with 2 degrees of freedom.

Zung (1997, 2000, 2003, 2006) gave (very mild) sufficient conditions on a compact  $r$ -dimensional orbit for the existence of a free system-preserving

Hamiltonian  $r$ -torus action near this orbit (hence, the Kalashnikov assumption about the existence of a circle action is not restrictive at all).

We extend the approach by Zung by considering a (so-called “hidden”) Hamiltonian torus action near a singular orbit, for real-analytic integrable systems. Such a torus action is generated by first integrals, some of which are multiplied with  $\sqrt{-1}$ . As we show, such an action is persistent under small real-analytic integrable perturbations of the system (hence, the assumption on the perturbed system to be semi-toric is not restrictive at all). We classified such torus actions up to symplectomorphisms, and proved their rigidity under perturbations.

By combining our approach and the third one (about a twist of the orbit), we extend the bifurcations of equilibria due to Duistermaat and van der Meer (with elliptic resonances) to compact orbits with twists. Using our approach of “hidden” torus actions, we construct two new series of singularities (with hyperbolic resonances and focus-focus resonances) that appear in generic integrable systems with 3 degrees of freedom.

## The velocity of one dimension cosmos

**Nenad Dj. Lazarov**

Department of Theoretical Physics and Condensed Matter Physics (020), Vinca  
Institute of Nuclear Sciences - National Institute of the Republic of Serbia,  
University of Belgrade, Republic of Serbia  
E-mail: [lazarov@vinca.rs](mailto:lazarov@vinca.rs)  
<https://orcid.org/0000-0003-3173-9636>

In this work we consider some properties which is generated by wellknown FRLW metric in the last hundred years . It will be presented geodesic equations in one space dimension. After solving geodesic equations in one space dimension and get the expression for velocity of tuba cosmos, the theoretical results will be compared with astronomical observation to estimate constant  $k$  which is parameter of geometric tensor in FRLW metric.

## A continuum space is the infinitely great

Qing Li

Shijiazhuang Traditional Chinese Hospital, China  
E-mail: liqingliyang@126.com

Regardless of how modern mathematics develops, it is impossible to calculate exact values (infinite values) with existing calculus because calculus are approximate. For example, for a given line segment or high-dimensional surface, we define it as a manifold that can be infinitely and arbitrarily divided into smaller parts. Here a certain line segment or surface is understood as a set of innumerable infinitesimals and can be calculated by calculus. From the new definition below, it can be seen that this above is not a fact. In this paper an infinitely small quantity is defined as a one-dimensional quantity of finite length without sizes of space, while an infinitely great quantity is reached by the superposition or accumulation of infinitely many finite quantities, by the way of the change in direction. The change in direction indicates that there is a jump from a finite quantity to infinitely many finite quantities (infinitely great). The form of the manifestation of the infinitely great is one quantitative continuum that cannot be operated by any algorithms, such as the operations of addition, subtraction, multiplication, and division. All parts of space we see is this one quantitative continuum that cannot be talked about anything outside of it and can compresses any quantities outside of it to nothing. The change in direction suggests that the infinitesimal are not parts of infinite great and it does not truly exist (it is only reference quantities that define the infinitely great) due to the infinitely great is only one quantitative continuum where there is only one quantity to exist. Therefore, any line segment or high-dimensional manifold we see in daily life are this one quantitative continuum. As a result from this new definition, the infinite exact value of a circumferential length ( $\pi$ ) has been obtained here.

## The Jacobi-orthogonality and Osserman tensors

Katarina Lukić

University of Belgrade, Faculty of Mathematics, Belgrade, Serbia

E-mail: katarina.lukic@matf.bg.ac.rs

<https://orcid.org/0000-0001-7638-8994>

We introduce a new potential characterization of Riemannian Osserman algebraic curvature tensors. An algebraic curvature tensor is Jacobi-orthogonal if  $\mathcal{J}_X Y \perp \mathcal{J}_Y X$  holds for all  $X \perp Y$ , where  $\mathcal{J}$  denotes the Jacobi operator. We prove that any Riemannian Jacobi-orthogonal tensor is Osserman, while all known Riemannian Osserman tensors are Jacobi-orthogonal. We generalize the property of Jacobi-orthogonality to indefinite scalar product spaces. We compare various principles and investigate relations between Osserman, Jacobi-dual, and Jacobi-orthogonal algebraic curvature tensors. We show that every quasi-Clifford tensor is Jacobi-orthogonal. We prove that a Jacobi-diagonalizable Jacobi-orthogonal tensor is Jacobi-dual whenever  $\mathcal{J}_X$  has no null eigenvectors for all nonnull  $X$ . We show that any algebraic curvature tensor of dimension 3 is Jacobi-orthogonal if and only if it is of constant sectional curvature. We prove that every 4-dimensional Jacobi-diagonalizable algebraic curvature tensor is Jacobi-orthogonal if and only if it is Osserman.

## The structure of dendrites and dynamics of continuous maps on them

Elena N. Makhrova

Lobachevsky State University of Nizhni Novgorod, Russia

E-mail: elena\_makhrova@inbox.ru

<https://orcid.org/0000-0002-5890-3064>

By *continuum* we mean a compact connected metric space. A *dendrite* is a locally connected continuum without subsets homeomorphic to a circle. Dendrites appear as Julia sets in complex dynamical systems (see [1]), as  $\omega$ -limit set of some dynamical systems (see, for example, [2], [3]).

Let  $X$  be a dendrite  $f : X \rightarrow X$  is a continuous map. The structure of dendrites is very complicated (see, for example, [4]). So properties of continuous maps depends on the structure of dendrites (see [5] – [8]). In the report we study the relationship between the structure of dendrites and the dynamics of continuous mappings on dendrites

The study was supported by the Russian Science Foundation grant ? 24-21-00242, <https://rscf.ru/project/24-21-00242/>

### References

- [1] H.-O. Peitgen, P. Richter. *The Beauty of Fractals*, Springer-Verlag, Heidelberg, (1986).
- [2] S.J. Agronsky, J.G. Ceder. *What sets can be  $\omega$ -limit sets in  $E^n$* , Real Analysis Exchange, **17** (1991-1992), 97-109.
- [3] F. Balibrea, J.L. García-Guirao. *Continua with empty interior as  $\omega$ -limit sets*, Applied General Topology, **6** (2005), 195-205.
- [4] K. Kuratowski. *Topology*, vol. 2, Academic Press, New York, (1968).
- [5] L.S. Efremova, E.N. Makhrova, *One- dimensional dynamical systems*, Russian Mathematical Surveys **76**:5 (2021), 821–881.
- [6] E.N. Makhrova. *Monotone maps on dendrites*, Discontinuity, Nonlinearity, and Complexity, **9**:4 (2020), 541-552.

- [7] E.N. Makhrova. *The structure of dendrites with the periodic point property*, Rus. Math. (Iz. VUZ), **55**:11 (2011), 33-37.
- [8] E.N. Makhrova. *Structure of dendrites admitting an existence of arc horseshoe*, Rus. Math. (Iz. VUZ), **59**:8 (2015), 52-61.

## Some curvature properties of quarter-symmetric metric connection

Miroslav Maksimović<sup>1</sup> and Milan Zlatanović<sup>2</sup>

<sup>1</sup>University of Priština in Kosovska Mitrovica, Faculty of Sciences and Mathematics, Kosovska Mitrovica, Serbia

E-mail: miroslav.maksimovic@pr.ac.rs

<https://orcid.org/0000-0002-8997-2812>

<sup>2</sup>University of Niš, Faculty of Science and Mathematics, Niš, Serbia

E-mail: milan.zlatanovic@pmf.edu.rs

<https://orcid.org/0000-0002-0318-1092>

In this talk, we will discuss the quarter-symmetric metric connection in generalized Riemannian manifolds. We will apply the obtained results to Kähler and cosymplectic (co- Kähler) manifolds, where we will determine tensors that do not depend on the generator of this connection and demonstrate their application in determining specific relations for the Weyl projective curvature tensor and the holomorphically projective curvature tensor.

The first author supported by the project of Faculty of Sciences and Mathematics, University of Priština in Kosovska Mitrovica (internal-junior project IJ-2303).

The second author acknowledge the grant of the Ministry of Education, Science and Technological Development of Serbia 451-03-65/2024-03/200124 for carrying out this research.



## **Ricci-Bourguignon Almost Solitons with Special Potential on Sasaki-like Almost Contact Complex Riemannian Manifolds**

**Mancho Manev**

Paisii Hilendarski University of Plovdiv & Medical University - Plovdiv, Bulgaria

E-mail: mmanev@uni-plovdiv.bg

<https://orcid.org/0000-0001-6181-8955>

Almost contact complex Riemannian manifolds, known also as almost contact B-metric manifolds, are equipped with a pair of pseudo-Riemannian metrics that are mutually associated to each other using the tensor structure. Here we consider a special class of these manifolds, those of the Sasaki-like type. They have an interesting geometric interpretation: the complex cone of such a manifold is a holomorphic complex Riemannian manifold (also called a Kähler-Norden manifold). The basic metric on the considered manifold is specialized here as a soliton, i.e. has an additional curvature property such that the metric is a self-similar solution of an intrinsic geometric flow. Almost solitons are more general objects than solitons because they use functions rather than constants as coefficients in the defining condition. A  $\beta$ -Ricci-Bourguignon-like almost soliton ( $\beta$  is a real constant) is defined using the pair of metrics. The introduced soliton is a generalization of some well-known (almost) solitons (such as those of Ricci, Schouten, Einstein), which in principle arise from a single metric rather than a pair of metrics. The soliton potential is chosen to be collinear to the Reeb vector field. The resulting manifolds equipped with the introduced almost solitons are characterized geometrically. In the three-dimensional case, an explicit example is constructed and the properties obtained in the theoretical part are confirmed.

## Lorentz Transformation and time dilatation

Miodrag Mateljević

Mathematical Institute, Serbian Academy of Sciences and Arts, Belgrade, Serbia

E-mail: miodrag@matf.bg.ac.rs

<https://orcid.org/0000-0002-9226-0023>

We consider two inertial frames  $S$  and  $S'$  and suppose that frame  $S'$  moves, for simplicity, in a single direction: the  $x$ -direction of frame  $S$  with a constant velocity  $v$  as measured in frame  $S$ .

Using homogeneity of space and time we derive modified Lorentz Transformation (LT) between two inertial reference frames without using the second postulate of Einstein, i.e., we do not assume the invariant speed of light (in vacuum) under LT.

Roughly speaking we suppose :

(H) Any clock which is at rest in its frame measures a small increment of time by some factor  $s = s(v)$ .

For  $s = 1$  we get the Galilean transformation of Newtonian physics, which assumes an absolute space and time. We also consider relation between absolute space and Special Relativity Theory, thereafter STR. As a corollary of relativity theory experimentally verified assumption is

(TD): (H) holds with Lorentz factor  $1/\gamma$ .

It seems here that we need physical explanation for (TD).

We show that (TD) is equivalent with

Postulate 3. The two-way speed is  $c$  in any inertial frame.

Note that Postulate 3 is a weaker assumption than Einstein second postulate.

We can derive the corresponding (LT) and explain twin paradox.

We introduce Postulate 3. The two- way speed of light in  $x'$  and  $z'$ -directions of the frame  $S'$  are  $c$ , and outline derivation of (LT) in this setting. Note that Postulate 3 is a weaker assumption than Einstein second postulate.

We also consider connection with hyperbolic and Minkowski geometry.

## Calculation of quiver representations of finite ring digraphs

Jelena Matović<sup>1</sup>, Aleksandar Lipkovski<sup>2</sup>

<sup>1</sup>Academy of Technical and Art Applied Studies - School of Electrical and  
Computer Engineering, Belgrade, Serbia  
jelena.matovic@viser.edu.rs

<sup>2</sup>University of Belgrade - Faculty of Mathematics, Belgrade, Serbia  
E-mail: aleksandar.lipkovski@matf.bg.ac.rs  
<https://orcid.org/0000-0002-7267-1490>

The authors have introduced finite ring digraphs [1] and recently studied the quiver point of view [2]. This is a continuation of this work, containing calculations of quiver representations which correspond to simplest types of these digraphs, in an attempt to introduce cohomological techniques in the topic.

### References

- [1] A. T. Lipkovski: *Strukturne grafi kolec: opredelenia i pervie rezuljtati*. Fundamentallnaja i prikladnaja matematika, 2015, tom 20, No 3, s. 181190.
- [2] A. T. Lipkovski, J. Matovic: *Quivers associated with finite rings - a cohomological approach*. Filomat 37:25 (2023), 85838589

## Geodesic mappings and their generalizations

Josef Mikes<sup>1</sup>, Nadezda Guseva<sup>2</sup>, Irena Hinterleitner<sup>3</sup>, Patrik Peška<sup>4</sup>, Lenka Vítková<sup>5</sup>

<sup>1</sup>Palacký University Olomouc, Czech Republic

E-mail: josef.mikes@upol.cz

<https://orcid.org/0000-0003-2098-7976>

<sup>2</sup>Moscow Pedagogical State University, Moscow, Russia

<sup>3</sup>Brno University of Technology: Brno, Czech Republic

<https://orcid.org/0000-0002-1046-578X>

<sup>4</sup>Palacký University Olomouc, Czech Republic

<https://orcid.org/0000-0002-0296-268X>

<sup>5</sup>Brno University of Technology: Brno, Czech Republic

<https://orcid.org/0000-0003-4805-908X>

A significant aspect of modern differential geometry revolves around the examination of mappings that transform geodesics into particular curves, such as geodesics, nearly geodesic curves, analytic curves, and so forth. The analysis of these mappings holds relevance in theoretical physics, particularly in the representation of gravitational, electromagnetic, and various other fields.

Geodesic mappings are mappings that map any geodesic onto geodesic. Let  $A_n = (M, \nabla)$  and  $\bar{A}_n = (\bar{M}, \bar{\nabla})$  be affine spaces, where  $M$  and  $\bar{M}$  are  $n$ -dimensional manifolds,  $\nabla$  and  $\bar{\nabla}$  are torsion-free affine connections. It is known [6] that  $A_n$  admits geodesic mapping onto  $\bar{A}_n$  if and only if the *Levi-Civita equations*

$$\bar{\nabla}_X Y = \nabla_X Y + \psi(X)Y + \psi(Y)X \quad (4)$$

hold for any tangent fields  $X, Y$  and where  $\psi$  is a linear differential form,  $\nabla$  and  $\bar{\nabla}$  are the Levi-Civita connections. We can suppose that  $\bar{M} = M$ . If  $\psi$  is vanishing then the mapping  $f$  is *affine* or *trivially geodesic*.

It has been demonstrated that the geodesic mappings of (pseudo-)Riemannian spaces onto (pseudo-)Riemannian spaces preserve differentiability class of the metric [3].

This preservation of differentiability class also extends to geodesic mappings from spaces endowed with affine connections to (pseudo-)Riemannian spaces [4].

Moreover, it has been established that all affine spaces are geodesically equivalent to equiaffine spaces [16]. Mikeš and Berezovski [7] devised a criterion, equivalent to the Levi-Civita equations, for determining the existence of geodesic mappings from equiaffine spaces to (pseudo-)Riemannian spaces. This criterion is formulated as a system of linear partial differential equations involving covariant derivatives. Importantly, these equations hold true both globally and locally, see [6]. A detailed look at geodesic mappings of two-dimensional spaces can be found in [5,13], and parabolic Kähler spaces in [11].

A specific focus is dedicated to the investigation of what is termed as geodesic bifurcation. This phenomenon arises when multiple geodesics go through a given point in a given direction, in the case where the condition of geodesic uniqueness is not fulfilled. An illustration of geodesic bifurcation was provided within the context of surfaces of revolution. Furthermore, building upon this illustration, additional examples were presented, including that of closed geodesics, as well as instances of geodesic bifurcation within product spaces [14].

Subsequently, S.G. Leiko introduced the concept of rotary mapping, which refers to mappings that map any geodesic into an isoperimetric extremal of rotation. An isoperimetric extremal of rotation is a distinctive curve that serves as an extremum for both the length functional and the rotation functional. These curves hold significance in physics as they represent the trajectories of particles with a spin. We extended this concept and found a counterexample of spaces that admit rotary mappings [8]. A detailed study of these questions is in our works [15,17].

Lastly, we touch upon the fundamentals of projective transformation and the theory of holomorphically projective transformations, particularly of Kähler manifolds. Additionally, we provide a brief overview of  $F$ -planar mappings, giving their definitions and presenting selected results derived from these theories [2,6,9,10,18].

## References

- [1] Ch. Almazbekov, N.I. Guseva, J. Mikeš, *Fundamental equations on conformal Fedosov spaces*. Springer Proc. Math. & Statistics 440, 223–229, 2024.
- [2] I. Hinterleitner, *Holomorphically projective mappings of (pseudo-) Kähler manifolds preserve the class of differentiability*. Filomat, 30:11, 3115–3122, 2016.

- [3] I. Hinterleitner, J. Mikeš, *Geodesic mappings and differentiability of metrics, affine and projective connections*. Filomat, 29:6, 1245–1249, 2015.
- [4] I. Hinterleitner, J. Mikeš, *Geodesic mappings onto Riemannian manifolds and differentiability*. Geometry, Integrability and Quantization, 18, 183–190, 2017.
- [5] I. Hinterleitner, N. Guseva, J. Mikeš, *On geodesic definiteness by similarity points*. J. Math. Sci (US) 277(5), 727–735, 2023.
- [6] J. Mikeš et al., *Differential Geometry of Special Mappings*. Palacky Univ. Press, Olomouc, 2019.
- [7] J. Mikeš, V. Berezovski, *Geodesic mappings of affine-connected spaces onto Riemannian spaces*. Amsterdam: North-Holland. Colloq. Math. Soc. J. Bolyai, 56. Diff. Geom. Eger, 491–494, 1989.
- [8] J. Mikeš, L. Rýparová, H. Chudá, *On the theory of rotary mappings*. Math. Notes 104:4, 617–620, 2018.
- [9] J. Mikeš, N.I. Guseva, P. Peška, L. Rýparová, *Rotary mappings and projections of a sphere*. Math. Notes 110:1, 152–155, 2021.
- [10] J. Mikeš, N.I. Guseva, P. Peška, L. Rýparová, *Almost geodesic mappings and projections of the sphere*. Math. Notes 111:3, 498–502, 2022.
- [11] R. Mohammad, M. Sheha, J. Mikeš, *On equidistant parabolic Kähler manifolds and geodesic mappings*. Filomat 38(3), 1033–1041, 2024.
- [12] P. Peška, M. Jukl, J. Mikeš, *Tensor decompositions and their properties*. Mathematics 11(17), N. 3638, 2023.
- [13] P. Peška, L. Vítková, J. Mikeš, I. Kuzmina, *On general solutions of Sinyukov equations on two-dimensional equidistant (pseudo-)Riemannian spaces*. Springer Proc. Math. & Statistics 440, 209–221, 2024.
- [14] L. Rýparová, J. Mikeš, A. Sabykanov, *On geodesic bifurcations of product spaces*. J. Math. Sci., New York 239:1, 86–91, 2019.
- [15] L. Rýparová, J. Mikeš, *Rotary mappings and rotary transformations*. J. Math. Sci., New York 276(4), 558–562, 2023.
- [16] J.M. Thomas, Asymmetric displacement of a vector. Trans. AMS 28:4, 658–670, 1926.
- [17] L. Vítková, *Rotary mappings of equidistant spaces*. Springer Proc. Math. & Statistics 440, 231–238, 2024.
- [18] L. Vítková, I. Hinterleitner, J. Mikeš, *Rigidity of holomorphically projective mappings of Kähler spaces with finite complete geodesics*. Mathematics 12(8), N. 1239, 2024.

## **New types of mappings of generalized Riemannian spaces**

**Vladislava Milenković**

Faculty of Technology, University of Niš, Serbia

E-mail: [vanja.dunja91@gmail.com](mailto:vanja.dunja91@gmail.com)

<https://orcid.org/0000-0003-0925-5133>

The talk deals with some new types of mappings of generalized Riemannian spaces in the Eisenhart sense. Some relations between corresponding curvature tensors of the generalized Riemannian spaces  $GR_N$  and  $G\bar{R}_N$  are presented, as well as some invariant geometric objects.

This work is financially supported by Serbian Ministry of Science, Technological Development and Innovation, Grant No. 451-03-65/2024-03/ 200133.



## Minimal models of nilmanifolds and complex structures

Dmitry Millionshchikov

Lomonosov Moscow State University and Steklov Institute of RAS, Russian Federation

E-mail: dmitry.millionschikov@math.msu.ru

<https://orcid.org/0000-0002-9149-467X>

Let  $G$  be a real nilpotent Lie group with the tangent Lie algebra  $\mathfrak{g}$  and  $J : \mathfrak{g} \rightarrow \mathfrak{g}$  define a left-invariant almost complex structure on  $G$ . The complexification  $\mathfrak{g}_{\mathbb{C}}^*$  of its dual space  $\mathfrak{g}^*$  splits to

$$(\mathfrak{g}^*)^{\mathbb{C}} = \Lambda^{1,0} \oplus \Lambda^{0,1}, \quad (5)$$

where  $\Lambda^{1,0}$  and  $\Lambda^{0,1}$  are eigen-spaces of  $J_{\mathbb{C}}^* : \mathfrak{g}^* \rightarrow \mathfrak{g}^*$  corresponding to  $\lambda = i, -i$ .  $J$  is integrable if and only if

$$d\Lambda^{1,0} \subset \Lambda^{1,0} \wedge (\Lambda^{1,0} \oplus \Lambda^{0,1}) = \Lambda^{2,0} \oplus \Lambda^{1,1},$$

where  $d : \mathfrak{g}_{\mathbb{C}}^* \rightarrow \mathfrak{g}_{\mathbb{C}}^* \wedge \mathfrak{g}_{\mathbb{C}}^*$  is dual to the Lie bracket  $[\cdot, \cdot]_{\mathbb{C}} : \mathfrak{g}_{\mathbb{C}} \wedge \mathfrak{g}_{\mathbb{C}} \rightarrow \mathfrak{g}_{\mathbb{C}}$ .

We define an increasing sequence of subspaces in the dual space  $\mathfrak{g}^*$

$$0 \subset V_1 \mathfrak{g}^* \subset V_2 \mathfrak{g}^* \subset \dots \subset V_l \mathfrak{g}^* \subset \dots \subset V_s \mathfrak{g}^* = \mathfrak{g}^*,$$

where  $V_l \mathfrak{g}^* = (\mathfrak{g}^{l+1})^{ann}$  is the annihilator of the ideal  $\mathfrak{g}^{l+1}$  of the lower central series. We define also the increasing sequence of holomorphic subspaces  $V_l^{1,0}, l \geq 1$ , of  $\Lambda^{1,0}$

$$V_l^{1,0} = (V_l \mathfrak{g}^*)^{\mathbb{C}} \cap \Lambda^{1,0}, \quad l = 1, \dots, s.$$

Successively expanding the bases of  $V_l^{1,0}$  Salamon [1] constructed a special basis  $\omega_1, \dots, \omega_n$  of the space  $\Lambda^{1,0}$  of holomorphic forms

$$d\omega^{l+1} \in I(\omega^1, \dots, \omega^l), \quad l = 0, \dots, n-1, \quad (6)$$

where  $I(\omega^1, \dots, \omega^l)$  is the ideal generated by forms  $\omega^1, \dots, \omega^l$  [1]. There are complex structures  $J$  (called nilpotent [2], [3]) for which the stronger condition of decomposability of  $d$  is satisfied

$$d\omega^{l+1} \in \Lambda^2(\omega^1, \dots, \omega^l, \bar{\omega}^1, \dots, \bar{\omega}^l), \quad l = 0, \dots, n-1, \quad (7)$$

The differential algebra generated by  $\omega^1, \dots, \omega^l$  can be considered as a special minimal model  $\mathcal{M}_{\mathfrak{g}}^J$  of a nilmanifold  $G/\Gamma$ , where  $\Gamma$  is a cocompact lattice. The existence of nilpotent complex structure  $J$  imposes restrictions on the structure of  $\mathfrak{g}$ . For example, an estimate for the first Betti number  $b_1(\mathfrak{g}) \geq 3$ . Recently, interest has grown in nilmanifolds, that admit only nonnilpotent complex structures. The first such example was found by Salamon [1]. In [3] Lie algebras with a one-dimensional center were studied, they also do not admit nilpotent complex structures.

We introduce the notion of a special minimal model  $\mathcal{M}_{\mathfrak{g}}^J$ , which is constructed from the nilpotent Lie algebra  $\mathfrak{g}$  taking into account the complex structure of  $J$ . With its help the theorem is proved.

**Theorem** A real 8-dimensional nilpotent Lie algebra  $\mathfrak{g}$  such that  $b_1(\mathfrak{g}) = 2$  admits an integrable complex structure if and only if  $\mathfrak{g}$  is isomorphic to one and only one Lie algebra from the following five Lie algebras

$$\mathcal{L}(2, 4), \mathfrak{n}_1^+(5)_0, \mathfrak{n}_1^+(5)_1, \mathfrak{g}_0, \mathfrak{g}_1.$$

In [1], [3] a short notation of commutation relations for nilpotent Lie algebras is used. With its help we write the relations of

- $\mathcal{L}(2, 4) = (0, 0, 12, 13, 23, 14 + 25, 14 - 25, 15 + 24);$
- $\mathfrak{n}_1^+(5)_t = (0, 0, 12, 13, 23, 14 + 25, 16 + 35 + t \cdot (14 - 25), -26 + 34 + t \cdot (15 - 24);$
- $\mathfrak{g}_t = (0, 0, 12, 13, 23, 14 + 25, 14 + 3 \cdot 25, 16 + 27 + t \cdot (14 + 25).$

Note also that the Lie algebras  $\mathfrak{n}_1^+(5)_t, t = 0, 1$ , are contained in the list [3] of 8-dimensional nilpotent Lie algebras with one-dimensional center and admitting a complex structure.

## References

- [1] S.M. Salamon, *Complex Structures on Nilpotent Lie Algebras*, J.Pure Appl.Algebra, **157** (2001), 311–333.

- [2] L.A. Cordero, M. Fernandez, A. Gray, L. Ugarte, *Compact nilmanifolds with nilpotent complex structures: Dolbeault cohomology*, Trans. Amer. Math. Soc. **352**:2 (2000), 5405–5433.
- [3] A. Latorre, L. Ugarte, R. Villacampa, *Complex structures on nilpotent Lie algebras with one-dimensional center*, J. of Algebra, **614** (2024), 271–306

## **Timelike Surfaces with Parallel Normalized Mean Curvature Vector Field and their Canonical Parameters**

**Velichka Milousheva**

Institute of Mathematics and Informatics, Bulgarian Academy of Sciences, Sofia,  
Bulgaria

E-mail: [vmil@math.bas.bg](mailto:vmil@math.bas.bg)

<https://orcid.org/0000-0001-6852-4749>

We introduce special isotropic parameters (which we call canonical parameters) for the class of timelike surfaces with parallel normalized mean curvature vector field in the Minkowski 4-space. These parameters allow us to describe this class of surfaces in terms of three geometrically determined functions. We prove a Fundamental existence and uniqueness theorem stating that each timelike surface with parallel normalized mean curvature vector field is determined up to a rigid motion in the Minkowski space by three geometric functions satisfying a system of three partial differential equations. In this way, we minimize the number of functions and the number of equations determining the surface, thus solving the Lund-Regge problem for this class of surfaces.

The talk is based on a joint work with Victoria Bencheva.

The author is partially supported by the National Science Fund, Ministry of Education and Science of Bulgaria under contract KP-06-N52/3.

## **On curvature tensors obtained by two non-symmetric linear connections**

**Svetislav M. Minčić**

Niš, Serbia

In the present paper, this problem is considered more generally by help of polylinear mappings and eight curvature tensor fields are obtained. Further, it is proved that among these fields five of them are independent, while the rest are linear combinations of the cited five fields.

## Maslov index and infinitesimal Lagrangian manifolds

Alexandr Mishchenko

Lomonosov Moscow State University, Russian Federation  
E-mail: asmish-prof@yandex.ru

The definition of the Maslov index of a Lagrangian manifold as a class of one-dimensional cohomology on it has given rise to numerous works generalizing the concepts of the Maslov index. In the works of V.I. Arnold [2,3], V.A. Vasiliev [4] and their followers, the theory of Lagrangian bordisms was developed and characteristic classes of Lagrangian submanifolds were constructed on its basis.

There is another approach to describing Maslov classes of Lagrangian submanifolds, presented in the works of V.V. Trofimov and A.T. Fomenko [5] from a categorical point of view.

In addition, there is a series of works by M.V. Karasev and V.P. Maslov [6] in which the canonical operator is constructed for general symplectic manifolds by covering a symplectic manifold with charts diffeomorphic to the simplest symplectic manifold.

Inspired by the works of V.V. Trofimov and A.T. Fomenko, we introduce the concept of so-called infinitesimal Lagrangian manifolds, which, in our opinion, allow us to characterize the characteristic classes of Lagrangian manifolds with maximum completeness and calculate the Maslov index for almost any Lagrangian manifolds.

The Maslov index is constructed as a homological invariant on a Lagrangian submanifold of some symplectic manifold. In the simplest case, the Lagrangian submanifold  $\Lambda \subset \mathbb{R}^{2n} \approx \mathbb{R}^n \oplus \mathbb{R}^n$  is a submanifold in the symplectic space  $\mathbb{R}^n \oplus \mathbb{R}^n$ , the symplectic structure in which is given by the non-degenerate form  $\omega = \sum_{i=1}^n dx^i \wedge dy^i$ , and  $\Lambda \subset \mathbb{R}^{2n}$  — is a submanifold,  $\dim \Lambda = n$ , on which the form  $\omega$  is trivial. In the general case, the symplectic manifold  $(W, \omega)$  is considered and a bundle of Lagrangian Grassmannians over a base parametrized by points of a symplectic manifold  $W$ .

The question that interests us is the following: when the Maslov index, given on an individual Lagrangian manifold as a one-dimensional cohomology class, is the image of some one-dimensional cohomology class of the total bundle space of Lagrangian Grassmannians. An answer is given for various classes of bundles of Lagrangian Grassmannians.

In particular, homological conditions are found for symplectic manifolds, under which explicit formulas are constructed for calculating the Maslov index of Lagrangian submanifolds.

The infinitesimal approach allows us to use the method of V.V.Trofimov again. Judging by his work (1994) [7], V.V.Trofimov was not satisfied with his generalization of Maslov classes, since with a large holonomy group, Maslov classes are lost. Here, too, we propose to formulate the problem in a categorical language. The object is a pair  $h : \Lambda \subset W$ , a Lagrangian submanifold  $\Lambda$  in a symplectic manifold  $W$ .

It is proposed to replace the symplectic manifold  $W$  with a smaller symplectic submanifold  $h' : \Lambda \subset W' \subset W$ . We get a wider set of characteristic classes, to which the old classes are reduced. Note that when defining characteristic classes, we use only the tangent bundle structure  $T(W)$  to a symplectic manifold  $W$ , a natural idea arises to replace the pair  $h : \Lambda \subset W$  on the Lagrangian manifold  $\Lambda$  itself, which is equipped with a Lagrangian structure using two vector bundles: tangent bundle of Lagrangian manifold and complex vector bundle with symplectic structure.

### **Conclusions.**

1. Every infinitesimal Lagrangian manifold can be realized as a Lagrangian submanifold of some symplectic manifold whose structure group of the tangent bundle is reduced to the subgroup  $\mathbb{O}(n)$ . This is enough to describe one-dimensional Maslov classes for an infinitesimal Lagrangian manifold.

2. The symplectic manifold from the previous paragraph can be chosen as the total space of the cotangent bundle of some  $n$ -dimensional manifold. It is unclear whether it is possible to embed an infinitesimal Lagrangian manifold into a compact symplectic manifold.

3. The problem of applying infinitesimal Lagrangian manifolds to asymptotic methods in mathematical physics equations is awaiting its solution.

## References

- [1] A. S. Mishchenko, *Maslov Index on Symplectic Manifolds. With Supplement by A. T. Fomenko "Constructing the Generalized Maslov Class for the Total Space  $W = T^*(M)$  of the Cotangent Bundle"*, Mathematical Notes, 112, 697708 (2022).
- [2] V. I. Arnol'd, *Lagrange and legendre cobordisms. I*, Functional Analysis and Its Applications, 14, 167177 (1980).
- [3] V. I. Arnol'd, *Lagrange and legendre cobordisms. II*, Functional Analysis and Its Applications, 14, 252260 (1980).
- [4] A. M. Vassiliev, *Lagrange and Legendre characteristic classes*, 2-d edition, Gordon and Breach Publishers, New York a.o., 1993, 273 pp.
- [5] V. V. Trofimov, A. T. Fomenko, *Algebra and geometry of integrable Hamiltonian differential equations*, *Mat. Prilozh.*[Mathematics and its Applications] Izdatelstvo "Faktorial", Moscow; copublished with Izdatelstvo Udmurtskogo Universiteta, Izhevsk, 1995. 448 pp.
- [6] M. V. Karasev, V. P. Maslov, *Pseudodifferential operators and a canonical operator in general symplectic manifolds*, Mathematics of the USSR-Izvestiya, Volume 23, Number 2, 277–305.
- [7] V. V. Trofimov, *Generalized Maslov classes on the path space of a symplectic manifold*. Trudy Mat. Inst. Steklov.205(1994), 172199; translation in Proc. Steklov Inst. Math.(1995), no.4, 157179.



## Weyl transform of a measure

Mansi Mishra

Indian Institute of Technology, B.H.U., India  
E-mail: mansimishra.rs.mat19@itbhu.ac.in

(1) Suppose  $\mu$  is a smooth measure on a smooth hypersurface of positive Gaussian curvature in  $R^{2n}$ .  $n \geq 2$ , then  $W(\mu)$ , the Weyl transform of  $\mu$ , is a compact operator, and if  $p > n \geq 6$ , then  $W(\mu)$  belongs to the  $p$ -Schatten class.

(2) There exist Schatten class operators with linearly dependent quantum translates.

## On deformations preserving dual arc length in dual 3-space

Marija Najdanović

University of Priština in Kosovska Mitrovica, Faculty of Sciences and Mathematics  
Kosovska Mitrovica, Serbia

E-mail: [marija.najdanovic@pr.ac.rs](mailto:marija.najdanovic@pr.ac.rs)

<https://orcid.org/0000-0003-4149-9664>

In this talk, we will discuss the geometrical aspects of dual curves, their properties and characterizations. Based on E. Study's mapping, the set of all oriented lines in Euclidean 3-space is in one to one correspondence with the set of points of dual unit sphere in dual 3-space and a differentiable curve on dual unit sphere represents a ruled surface. We will define a deformation of dual curves preserving dual arc length and consider some geometrical properties under this type of deformation. Consequently, observing the deformations of dual spherical curves we will actually study the deformations of the corresponding ruled surfaces. This is a joint work with Ljubica Velimirović and Svetozar Rančić.

*Acknowledgement.* This talk is supported by the project IJ2303 of Faculty of Sciences and Mathematics, University of Priština in Kosovska Mitrovica.

## On the stability of the identity map of compact space forms

Crina-Daniela Neacșu<sup>1</sup>, Gabriel-Eduard Vilcu<sup>2</sup>

<sup>1</sup>National University of Science and Technology Politehnica Bucharest, Romania

E-mail: crina.neacsu@upb.ro

<sup>2</sup>University of Bucharest & National University of Science and Technology

Politehnica Bucharest, Romania

E-mail: gabriel.vilcu@upb.ro

In this paper, we report on some recent results concerning the stability of the identity map on compact locally conformal almost cosymplectic manifolds of pointwise constant  $\phi$ -holomorphic sectional curvature and compact  $T$ -space forms.

## On geodesic orbit nilmanifolds

**Yurii Nikonorov**

Southern Mathematical Institute of the Vladikavkaz Scientific Center of the  
Russian Academy of Sciences, Russian Federation  
E-mail: nikonorov2006@mail.ru

The talk is devoted to recent results on geodesic orbit Riemannian metrics on nilpotent Lie groups (geodesic orbit nilmanifolds). In particular, we discuss the construction of continuous families of pairwise non-isomorphic connected and simply connected nilpotent Lie groups, every of which admits geodesic orbit metrics. The minimum dimension of such groups is 10.

## **Killing tensors on symmetric spaces**

**Yuri Nikolayevsky**

La Trobe University, Department of Mathematical and Physical Sciences,  
Australia

E-mail: [Y.Nikolayevsky@latrobe.edu.au](mailto:Y.Nikolayevsky@latrobe.edu.au)

<https://orcid.org/0000-0002-9528-1882>

I will discuss some known and new results, including a surprising construction showing that the algebra of Killing tensors on a Riemannian symmetric space is not in general generated by Killing vectors. This is a joint work with V. Matveev (Jena, Germany).

## **Implementation of Postprocessing Procedure of a Rapid Algorithm of Geometric Coding of Digital Images Using CUDA Architecture**

**Gleb Nosovski**

Faculty of Mechanics and Mathematics, Lomonosov Moscow State University,  
Moscow, Russian Federation

E-mail: [gleb.nosovski@gmail.com](mailto:gleb.nosovski@gmail.com), [gleb.nosovski@math.msu.ru](mailto:gleb.nosovski@math.msu.ru)

The talk presents some improvements of fast contour recognition algorithm. The main idea of this algorithm is based on calculation of geometric characteristics of a two-dimensional surface in  $R^3$ , which encodes digital images. To obtain the contours, Minkowski fractal dimension calculation is used. A postprocessing algorithm of contour thinning without loss of quality of entire contour picture is proposed in the paper. An estimate of the speed of the new algorithm is presented in comparison with the well-known implementation of the Canny algorithm using the OpenCV computer vision library and the parallel architecture of CUDA. Examples demonstrating the efficiency of the new algorithm are presented.

This is a joint work with A.Yu Chekunov.

## Fischer-Marsden Equation on Paracontact Geometry

Mustafa Özkan<sup>1</sup> and İrem Küpeli Erken<sup>2</sup>

<sup>1,2</sup>Bursa Technical University, Faculty of Engineering and Natural Science,  
Department of Mathematics, Bursa, Turkey

<sup>1</sup>E-mail: mustafa.ozkan@btu.edu.tr

<https://orcid.org/0000-0002-4483-2912>

<sup>2</sup>E-mail: irem.erken@btu.edu.tr

<https://orcid.org/0000-0003-4471-3291>

Let  $(M^{2n+1}, g)$  be a compact, orientable semi-Riemannian manifold. We denote the set of all unit volume semi-Riemannian metrics on  $(M^{2n+1}, g)$  by  $\mathcal{M}$ . The linearization of the scalar curvature  $\mathcal{L}_g(g^*)$  is given by

$$\mathcal{L}_g g^* = -\Delta_g(\text{tr}_g g^*) + \text{div}(\text{div}(g^*)) - g(g^*, S_g),$$

where  $\Delta_g$ ,  $\text{div}$ ,  $g^*$  and  $S_g$  denotes the negative Laplacian of the semi-Riemannian metric  $g$ , divergence operator, symmetric  $(0, 2)$  tensor field on  $M$  and the Ricci tensor, resp. The formal  $L^2$ -adjoint  $\mathcal{L}_g^* g^*$  of the linearized scalar curvature operator  $\mathcal{L}_g$  is defined by

$$\mathcal{L}_g^*(\lambda) = -(\Delta_g \lambda)g + \text{Hess}_g \lambda - \lambda S_g, \quad (8)$$

where  $\text{Hess}_g \lambda(X, Y) = \nabla_g^2 \lambda(X, Y) = g(\nabla_X D\lambda, Y)$  is the Hessian operator of the smooth function  $\lambda$  on  $M$  and  $D$  is the gradient operator of  $g$ . We refer the equation  $\mathcal{L}_g^*(\lambda) = 0$  as Fischer-Marsden equation (FME).

The goal of this study is to investigate the non-trivial solutions of Fischer-Marsden equation on paracontact geometry.

## **Characterizations of Ricci-Bourguignon almost solitons**

**Dhriti Sundar Patra**

Department of Mathematics, IIT Hyderabad Kandi, Sangareddy, India

E-mail: [dhriti@math.iith.ac.in](mailto:dhriti@math.iith.ac.in)

<https://orcid.org/0000-0002-7958-1690>

We characterize RB almost solitons on contact metric manifolds and also find some sufficient conditions under which these are isometric to a unit sphere or trivial (Einstein).



## **Integrability of the sub-Riemannian geodesic flow of the left-invariant metric on the Heisenberg group**

**Milan Pavlović, Tijana Šukilović**

University of Belgrade, Faculty of Mathematics, Belgrade, Serbia  
E-mail: milan.pavlovic@matf.bg.ac.rs, tijana.sukilovic@matf.bg.ac.rs

We study two different classes of normal geodesic flows corresponding to the left-invariant sub-Riemannian metric on the  $(2n + 1)$ -dimensional Heisenberg group. The first class corresponds to the left-invariant distribution, while the second corresponds to the right-invariant one. We show that corresponding Hamiltonian L-L and L-R systems are completely integrable.

## **Li-Yau sub-gradient estimates and Perelman-type entropy formulas for the heat equation in quaternionic contact geometry**

**Alexander Petkov**

Sofia University, Faculty of Mathematics and Informatics, Sofia, Bulgaria  
E-mail: a\_petkov\_fmi@abv.bg

We are going to present in this talk two sub-gradient estimates for the quaternionic contact (qc) heat equation on a compact qc manifold of dimension  $4n + 3$ , provided some positivity conditions are satisfied. These are qc versions of the prominent Li-Yau gradient estimate in Riemannian geometry. Another goal of the talk is to demonstrate two Perelman-type entropy formulas for the qc heat equation on a compact qc-Einstein manifold of dimension  $4n + 3$  with non-negative qc scalar curvature (e.g. compact 3-Sasakian manifold), as well as an integral sub-gradient estimate for the positive solutions of the qc heat equation.

The research is partially supported by National Science Fund of Bulgaria, National Scientific Program "VIHREN", Project KP-06-DV-7 and Contract KP-06-H72-1/05.12.2023 with the National Science Fund of Bulgaria.

## Composition of conformal and projective mappings of generalized Riemannian spaces in Eisenhart's sense preserving certain tensors

Miloš Petrović

University of Niš, Faculty of Agriculture in Kruševac, Kosančićeva 4, 37000  
Kruševac, Republic of Serbia  
E-mail: petrovic.milos@ni.ac.rs  
<https://orcid.org/0000-0002-0308-8074>

Conformal and projective mappings have important physical interpretations in space-times. These mappings have been thoroughly studied. Their composition between Riemannian spaces that was at the same time harmonic had been initiated by S. E. Stepanov, I. G. Shandra, Seven Classes of Harmonic Diffeomorphisms, Mathematical Notes, 2003, Volume 74, Issue 5, Pages 708-716 and further studied by I. Hinterleitner, Special mappings of equidistant spaces, Ph.D. Thesis, Brno University of Technology, Brno, 2009. Conformal and projective mappings of Riemannian spaces preserving certain tensors were studied by O. Chepurna, Diffeomorphisms of Riemannian spaces with preserved Einstein tensor, Ph.D. thesis, Faculty of Science, Palacky University, Olomouc, 2012. We consider conformal and projective mappings of generalized Riemannian spaces in Eisenhart's sense and find necessary and sufficient conditions for these mappings to preserve curvature, Ricci and Einstein tensors and some of their linear combinations. Particularly, as an additional contribution to related results collected in the Ph.D. thesis by O. Chepurna, we find that the following result holds in the case of Riemannian spaces: if a conformal mapping  $f_1 : M \rightarrow \hat{M}$  is preserving the Einstein tensor and a projective mapping  $f_2 : \hat{M} \rightarrow \overline{M}$  is preserving the Einstein tensor then the Yano tensor of concircular curvature is invariant with respect to the composition  $f_3 = f_1 \circ f_2 : M \rightarrow \overline{M}$ .

Research was financed by the Ministry of Science, Technological Development and Innovation, Republic of Serbia (Contract reg. no.451-03-65/2024-03/200383)

## Numerical algorithms on Hadamard manifolds

**Ariana Pitea**

National University of Science and Technology Politehnica Bucharest, Romania

E-mail: [arianapitea@yahoo.com](mailto:arianapitea@yahoo.com)

<https://orcid.org/0000-0001-8198-3077>

Numerical procedures for the reckoning of fixed points are analysed from a qualitative point of view in the geometric setting offered by Hadamard manifolds. Convergence, stability and data dependence results are presented, including applications.

## On certain new results on the Steenrod algebra $\text{mod } p$

**Theodore Popelensky**

Moscow State University, Moscow, Russia  
E-mail: popelens@gmail.com

For the Steenrod algebra  $\text{mod } p$ ,  $p > 2$ , three additive bases are well known. Namely, basis of admissible monomials, Milnor basis, and  $P_s^t$ -basis. To be more precise, there is a family of  $P_s^t$ -bases. We develop certain new additive bases in the Steenrod algebra  $\text{mod } p$ . We investigate which pairs of bases have triangular transition matrices. Finally we plan to discuss certain applications. If time permits we shall discuss certain generalization to Brown-Peterson cohomology.

## **The geometrical personalization of human organs 3D models by using the Characteristic Product Features methodology**

**Ljiljana Radović**

Faculty of Mechanical Engineering, University of Nis, Serbia  
E-mail: ljiljana.radovic@masfak.ni.ac.rs

Computer-assisted orthopedic surgery (CAOS) involves applying various computer-based methodologies and devices to plan, guide, and perform surgical procedures, thereby improving outcomes throughout the surgical process. This study integrates the Characteristic Product Features (CPFs) methodology developed in-house to improve CAOS. It enables the creation of the human organs' geometrical models by including different relations between Regions of Interest (RGIs) models and specific properties, like functional, materials, and topological. Enhancing existing methodologies in CAOS aims to offer a more comprehensive geometrical description of human organs, leading to the development of more precise and anatomically accurate personalized geometrical models. Creating customized geometry with accurately defined features is expected to enable surgeons to prepare and execute surgical interventions better, consequently improving patient care and recovery prospects. The demonstration of successful geometry adaptation will be shown by prototyping developed models using 3D FDM printing.

This is a joint work with Nikola Vitković, Ljiljana Radović, Jelena Stojković, Aleksandar Miltenović

## Basins of attractions of a new iterative method for finding simple zeros

Lidija Rančić<sup>1</sup> and Svetozar Rančić<sup>2</sup>

<sup>1</sup>University of Niš, Faculty of Electronic Engineering Niš, Serbia

E-mail: lidija.rancic@elfak.ni.ac.rs

<https://orcid.org/0000-0003-4888-4485>

University of Niš, Faculty of Sciences and Mathematics, Department of Computer Science, Niš, Serbia

E-mail: rancicsv@yahoo.com

<https://orcid.org/0000-0002-1023-3807>

We present new fourth order iterative method for solving nonlinear equations. New method is obtained by composition of super Halley's and Homer's three-order methods. Frequently, comparisons of the various schemes are based on the number of iterations required for convergence, number of function evaluations, and/or amount of CPU time. In addition to these criteria, we consider criteria based on basins of attraction. The basin of attraction is a method to visually comprehend how an algorithm behaves as a function of the various initial values. Through many numerical examples it is shown that new method can be competitive to other fourth-order methods.

*AMS(MOS) subject classification:* 65H05

This work was supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia [grant numbers 451-03-65/2024-03/200102 and 451-03-47/2023-01/200124]



## The Brylinski beta function of a double layer

Pooja Rani and M.K. Vemuri

Department of Mathematical Sciences, IIT (BHU), Varanasi, India  
E-mail: poojarani.rs.mat19@itbhu.ac.in, mkvemuri.mat@itbhu.ac.in

An analogue of Brylinski's knot beta function is defined for a compactly supported (Schwartz) distribution  $T$  on  $d$ -dimensional Euclidean space. This is a holomorphic function on a right half-plane. If  $T$  is a (uniform) double-layer on a compact smooth hypersurface, then the beta function has an analytic continuation to the complex plane as a meromorphic function, and the residues are integrals of invariants of the second fundamental form. The first few residues are computed when  $d = 2$  and  $d = 3$ .

## Weak nearly Sasakian and weak nearly cosymplectic manifolds

Vladimir Rovenski

Department of Mathematics, University of Haifa, Israel

E-mail: vrovenski@univ.haifa.ac.il

<https://orcid.org/0000-0003-0591-8307>

Nearly Kähler structure  $(J, g)$  was defined by A. Gray (1970) using the condition that the symmetric part of  $\nabla J$  vanishes. Nearly Sasakian and nearly cosymplectic structures were defined by D. Blair and his collaborators (1976) as odd- dimensional analogs of nearly Kähler structure. Nicola A., Dileo G. and Yudin I. [1] proved that nearly cosymplectic manifolds are Riemannian products of two kinds, and that nearly Sasakian manifolds of dimension greater than 5 are Sasakian. Weak contact manifolds, see [2], i.e., a nonsingular skew-symmetric tensor instead of the complex structure on the contact distribution, allow us to expand the theory of almost contact metric manifolds. Weak nearly Sasakian and weak nearly cosymplectic structures  $(\varphi, Q, \xi, \eta, g)$  were defined in [3] similarly to classical case (when  $Q = \text{Id}$ ) using certain conditions for the symmetric part of  $\nabla \varphi$ .

In this talk, we present theorems generalizing results of [1]: splitting of weak almost cosymplectic manifolds [4], and characterization of Sasakian manifolds in the class of weak contact metric manifolds [5,6].

### References

- [1] A.D. Nicola, G. Dileo, and I. Yudin, *On nearly Sasakian and nearly cosymplectic manifolds*, Annali di Matematica, 197 (2018), 127–138
- [2] V. Rovenski and R. Wolak, *New metric structures on  $\mathfrak{g}$ -foliations*, Indagationes Mathematicae, 33 (2022), 518–532
- [3] V. Rovenski, *Weak nearly Sasakian and weak nearly cosymplectic manifolds*. Mathematics 2023, 11(20), 4377
- [4] V. Rovenski, *On the splitting of weak nearly cosymplectic manifolds*. Preprint, 16 pp. arXiv:2310.19518, Accepted in DGA

- [5] V. Rovenski, *Foliated structure of weak nearly Sasakian manifolds*. Preprint, 10 pp. arXiv:2312.11411, Accepted in AMPA
- [6] V. Rovenski, *Characterization of Sasakian manifolds*. Preprint, 14 pp. arXiv:2401.02752, Accepted in AEJM

## Willmore-type variational problem for foliated hypersurfaces

Vladimir Rovenski

Department of Mathematics, University of Haifa, Israel

E-mail: vrovenski@univ.haifa.ac.il

<https://orcid.org/0000-0003-0591-8307>

After T. Willmore [4], many authors, including R. Reilly [2] and Z. Guo [1], searched for immersions of a manifold  $M$  in a Riemannian manifold, which are critical points of functionals  $W = \int_M H^p dV$  or  $J = \int_M \|h\|^p dV$ . Here,  $h$  is the 2nd fundamental form,  $H$  the mean curvature and  $p > 0$ . An interesting problem is the generalization of the Willmore functional to submanifolds with additional structures, such as almost products or foliations.

In the talk, we discuss critical points of the new Willmore-Reilly type functional  $WF = \int_M F(\sigma_1^{\mathcal{F}}, \dots, \sigma_s^{\mathcal{F}}) dV$  for a hypersurface  $M$  in  $\mathbb{R}^{n+1}$  equipped with an  $s$ -dimensional ( $s \leq n$ ) foliation  $\mathcal{F}$ , see [3]. Here,  $F \in C^3(\mathbb{R}^n)$  and  $\sigma_i^{\mathcal{F}}$  are elementary symmetric functions of the eigenvalues of  $h$  restricted on the leaves of  $\mathcal{F}$ . The 1st and 2nd variations of such functionals are calculated and the conformal invariance of some of them (type Guo) is proven. The corresponding Euler-Lagrange equations for a critical hypersurface with transverse harmonic foliation are found and examples with  $s \leq 2$  are given. Critical hypersurfaces of revolution in  $\mathbb{R}^{n+1}$  are presented and shown to be a local minimum for special variations of immersion.

### References

- [1] Z. Guo, *Generalized Willmore functionals and related variational problems*. Diff. Geom. Appl., 25 (2007), 543–551.
- [2] R.C. Reilly, *Variational properties of functions of the mean curvatures for hypersurfaces in space forms*. J. Differ. Geom. 8, 465–477 (1973).

- [3] V. Rovenski, Willmore-type variational problem for foliated hypersurfaces, preprint, 2024, 14 pages. arXiv.2402.17565.
- [4] T.J. Willmore, *Note on embedded surfaces*. An. Stiint. Univ. "Al. I. Cuza" Iasi, Sect. Ia Mat. 11. (1965), 493–496.

## On the Index Problem On Manifolds with Group Actions. Contributions of Fixed Points

Anton Savin

RUDN University, Moscow, Russia

E-mail: a.yu.savin@gmail.com

<https://orcid.org/0000-0002-7094-4117>

Given an action of a discrete group  $G$  on a smooth closed manifold  $X$ , we consider a representation of  $G$  by shift operators  $T_g : u(x) \mapsto u(g^{-1}x)$  in function spaces on  $X$  and a class of matrix operators with shifts of the form

$$D = \sum_{g \in G} D_g T_g : C^\infty(X, \mathbb{C}^N) \longrightarrow C^\infty(X, \mathbb{C}^N),$$

where the coefficients  $D_g$  are matrix (pseudo)differential operators on  $X$ . If appropriate ellipticity conditions are satisfied,  $D$  is a Fredholm operator and the problem arises of calculating its index in terms of the principal symbol  $\sigma(D)$  of this operator (see, for example, [1] and the cited literature). It should be noted that the principal symbol is an element of an essentially non-commutative algebra — the crossed product of the algebra of matrix functions on the cospheric bundle of the manifold and the group  $G$  acting on the indicated algebra by automorphisms.

In [2], the index was calculated in the case of the group  $G = \mathbb{Z}$  and it was shown that the index is equal to the pairing of the class in the  $K$ -theory of the crossed product defined by the principal symbol  $\sigma(D)$  and the equivariant Todd class of the manifold in periodic cyclic cohomology of the crossed product.

We define the Todd class for groups of the form  $G = \mathbb{Z} \oplus F$ , where  $F$  is a finite group, and prove the corresponding index formula. The main difficulties that had to be overcome: 1) the action of the group, generally speaking, is not assumed to be isometric; 2) it was necessary to describe the contributions of the fixed points of the group action to the index formula.

The results are obtained in a joint work with Haydar Abbas (RUDN University).

## References

- [1] A. Antonevich, A. Lebedev, *Functional-differential equations. I.  $C^*$ -theory*. Pitman Monographs and Surveys in Pure and Applied Mathematics, 70. — Harlow: Longman Scientific & Technical, 1994.
- [2] A. Savin, B. Sternin, *Index of elliptic operators for diffeomorphisms of manifolds* J. Noncommut. Geom., **83**:3 (2014), 695-734.

## Geometric Properties of the Inertia Tensor of a Rigid Body

Anastasiia Shubert

Moscow State University, Moscow, Russian Federation  
E-mail: anastasiia.shubert@math.msu.ru

The report will explore the connection between the following mechanical problems:

- the problem of the rotation of a rigid body around a fixed point (i.e., a spinning top) in a 3-dimensional (pseudo-)Euclidean space,
- the problem of the motion of a rigid body (referred to as a “plate”) in a 2-dimensional space of constant curvature, specifically, on a 2-dimensional sphere, a Euclidean plane, or a Lobachevsky plane.

The inertia tensor of the rigid body from mechanics will be examined. We will describe its connection with the kinetic energy of the rigid body and the inertia tensor on the Lie algebras  $so(3)$  and  $so(2,1)$  in terms of a natural isomorphism between this Lie algebra and the ambient (pseudo-)Euclidean space. We will compute the inertia tensor of any single-point body in terms of the (pseudo-)Euclidean metric of the ambient space. As a consequence, firstly, it follows that the value of the inertia tensor (as a quadratic form) on any time-like vector is non-negative. In particular, the inertia tensor cannot have a signature of  $(-, -, -)$ . Secondly, for any “plate” on the Lobachevsky plane (lying within the light cone), the inertia tensor is positive definite. Thirdly, we will provide specific examples of two-point bodies lying outside the light cone, whose inertia tensor can have any signature other than  $(-, -, -)$ .



## Invariants for geometric mappings

Dušan Simjanović<sup>1</sup>, Nenad Vesić<sup>2</sup>

<sup>1</sup>Metropolitan University, Belgrade, Serbia

E-mail: dsimce@gmail.com

<https://orcid.org/0000-0002-1709-0765>

<sup>2</sup>Mathematical Institute of Serbian Academy of Sciences and Arts, Serbia

E-mail: vesic.specijalac@gmail.com

<https://orcid.org/0000-0001-7070-7325>

In this research, we consider transformations of symmetric affine connections and invariants caused by them. Our results are classified in three sections. The first section is Introduction, where we recalled the basic definitions and motivations necessary for the research. In the second section, we considered transformations of affine connections generated with respect to metric tensors. In this section, we obtained the invariant geometrical objects with respect to the transformations of affine connections and curvature tensors of symmetric affine connection spaces with metrics defined on them. In the third section, we presented an example of application of results presented in the second section.

**Key words:** Riemannian space, sense, invariant

**Math. Subj. Classification:** 53A55, 53B20

<sup>2</sup>This research is financially supported by Serbian Ministry of Science, Technological Development, and Patents, through the Mathematical Institute of Serbian Academy of Sciences and Arts

## On some generalizations of Kahlerian spaces

Mića S. Stanković

University of Niš, Faculty of Science and Mathematics, Niš, Serbia

E-mail: mica.stankovic@pmf.edu.rs

<https://orcid.org/0000-0002-5632-0041>

In this talk, we present results on generalized Kahlerian spaces [1], generalized Kahlerian spaces of the first kind [2], and generalized Kahlerian spaces of the second kind [3]. In all three cases, invariant geometric objects of holomorphic projective mappings of generalized Kahlerian spaces, of the Weyl tensor type, are considered.

The author acknowledge the grant of the Ministry of Education, Science and Technological Development of Serbia 451-03-65/2024-03/200124 for carrying out this research.

### References

- [1] M. S. Stankovic, S. M. Mincic, Lj. S. Velimirovic, *On equitortion holomorphically projective mappings of generalised Kahlerian spaces*, Czechoslovak Mathematical Journal, vol.54 (129) , No. 3 (2004) 701-715.
- [2] M. S. Stankovic, M. Lj. Zlatanovic, Lj. S. Velimirovic, *Equitortion holomorphically projective mappings of generalized Kahlerian space of the first kind*, Czechoslovak Mathematical Journal, vol.Vol. 60, No. 3 (2010) 635-653.
- [3] M. S. Stankovic, M. Lj. Zlatanovic, L. S. Velimirovic, *Equitortion holomorphically projective mappings of generalized Kahlerian space of the second kind*, International Electronic Journal of Geometry, (M50) vol.Vol. 3, No. 2 (2010) 26-39.

## New vanishing theorem of conformal Killing forms

Sergey E. Stepanov

Finance University under the Government of the Russian Federation: Moscow,  
Russia

E-mail: s.e.stepanov@mail.ru

<https://orcid.org/0000-0003-1734-8874>

In this talk we generalize our result (see [1]) and results of Kashiwada and Kora (see [2] and [3]) in the form of the following theorem.

Theorem. Let  $(M, g)$  be an  $n$ -dimensional closed connected Riemannian manifold with non-positive curvature operator of the second kind. Then either it is a compact Euclidean space form, or it does not admit conformal Killing  $p$ -forms for all  $p = 1, 2, \dots, n-1$ .

### References

- [1] S.E. Stepanov, I.I. Tsyganok, *Theorems of existence and non-existence of conformal Killing forms*, Russian Math. (Iz. VUZ), 58:10 (2014), 4651
- [2] T. Kashiwada, *On conformal Killing tensor*, Natural Science Report, Ochanomizu University, 19:2 (1968), 67-74.
- [3] M. Kora, *On conformal Killing forms and the proper space of  $\Delta$  for  $p$ -forms*, Mathematical Journal of Okayama University, 22:2 (1980), 195-204.

## **Geometrical eigenproblem of various types higher order tensors in Riemannian space**

**Jelena Stojanov**

University of Novi Sad, Technical faculty "Mihajlo Pupin" Zrenjanin, Serbia

E-mail: [jelena.stojanov@uns.ac.rs](mailto:jelena.stojanov@uns.ac.rs)

<https://orcid.org/0000-0002-2237-8026>

The geometrical eigenproblem (G-eigenproblem) is validated as an extension of the classical eigenproblem of a linear operator within Euclidean vector space. It is homogeneous and invariant, making it efficient and advantageous for tensors describing real phenomena, regardless of the representation, while determining their eigenspaces. The G-eigenproblem in Riemannian space will be explored not only for covariant tensors but also for mixed and contravariant ones. Achievements will be discussed and related to higher-order unit tensors.

## **3-Triangulation of polyhedra and their connection graphs**

**Milica Stojanović**

University of Belgrade, Faculty of Organizational Sciences, Srbija

E-mail: milica.stojanovic@fon.bg.ac.rs

<https://orcid.org/0000-0002-2517-2189>

Here we investigate the properties of 3-triangulation of polyhedra, when possible. Namely, it is known that 3- triangulation of convex polyhedra is always possible, but this is not the case with all non-convex ones. This is the reason to consider the decomposition of non-convex polyhedra into convex pieces if possible. After that, we introduce the connection graph for the 3-triangulable polyhedron in such a way that these pieces are represented by the nodes of the graph.

First, our attention will be focused to the minimal number of tetrahedra necessary to 3-triangulate toroids, a special class of non-convex polyhedra. Polyhedra that are topologically equivalent to ball with  $p$  handles ( $p \in \mathbb{N}$ ) are called  $p$ -toroids, and more generally, independently on  $p$ , we call them toroids.

As another application of connection graphs, we shall also consider those corresponding to convex polyhedra, especially to conic triangulation of them.

## **Gromov-Hausdorff Distance and Geometric Optimization Problems**

**Alexey Tuzhilin**

Lomonosov Moscow State University, Moscow, Russian Federation

E-mail: tuz@mech.math.msu.su

We will discuss how the Gromov-Hausdorff distance arises in various combinatorial and geometric optimization problems. First, we recall our results (obtained jointly with Professor Alexander O. Ivanov) on (1) Borsuk's problem about the smallest number of parts into which a bounded subset of Euclidean space can be cut so that all parts have a smaller diameter than the original subset; (2) finding the smallest number of colors necessary to correctly color the vertices of the graph; (3) calculating the smallest number of cliques whose vertices cover the set of vertices of the graph; (4) calculating the weights of the edges of the minimum spanning tree in the weighted graph. After this, we will show how to express through the Gromov-Hausdorff distance the  $l_1$ -dimension, i.e., the smallest dimension of a vector space with Manhattan ( $l_1$ -) norm into which a given finite metric space can be isometrically embedded.

## Size in contact geometry

Igor Uljarević

University of Belgrade, Faculty of Mathematics

E-mail: [igor.uljarevic@matf.bg.ac.rs](mailto:igor.uljarevic@matf.bg.ac.rs)

<https://orcid.org/0000-0002-9405-1903>

In this talk, I will cover the basics of contact manifolds - defining them and highlighting their main properties. I will discuss the concept of contact non-squeezing, a handy tool for understanding size when there is no natural measure or metric. Lastly, I will share an interesting finding: while contact geometry does not capture size of a ball on a small scale, it surprisingly does so on a larger scale in certain cases.

## Extremal Kähler metrics and the moment map

Craig van Coevering

Bogazici University Istanbul, Turkey  
E-mail: craig.coevering@bogazici.edu.tr

An extremal Kähler metric is a canonical Kähler metric, introduced by E. Calabi, which is somewhat more general than a constant scalar curvature Kähler metric. The existence of such a metric is an ongoing research subject and expected to be equivalent to some form of geometric stability of the underlying polarized complex manifold  $(M, J, [\omega])$  –the *Yau-Tian-Donaldson conjecture*. Thus it is no surprise that there is a moment map, the scalar curvature (A. Fujiki, S. Donaldson), and the problem can be described as an infinite dimensional version of the familiar finite dimensional G.I.T.

In this talk I will describe how the moment map can be used to describe the local deformation problem of constant scalar curvature and extremal metrics. Essentially, the local picture can be reduced to finite dimensional G.I.T. In particular, we can construct a coarse moduli space of extremal Kähler metrics with a fixed polarization  $[\omega] \in H^2(M, \mathbb{R})$ , which is an Hausdorff complex analytic space.



## Conformal transformations of curvature pseudotensors

Ana Velimirović<sup>1</sup>, Milan Zlatanović<sup>2</sup>,

<sup>1</sup>University Metropolitan, Belgrade, Serbia

E-mail: velimirovic018@gmail.com

<https://orcid.org/0000-0002-2594-6615>

<sup>2</sup>Faculty of Science and Mathematics, University of Niš, Serbia

E-mail: milan.zlatanovic@pmf.edu.rs

<https://orcid.org/0000-0002-0318-1092>

In this talk we study curvature pseudotensors. They are tensors constructed from the curvature tensor and other geometric quantities. However, they are not true tensors since they don't transform like tensors under arbitrary coordinate transformations. Instead, they transform according to certain rules under specific transformations, such as conformal transformations. We exam ET conformal transformations at  $GR_N$ .

## Curvature based shape consideration

Ljubica Velimirović

Faculty of Science and Mathematics, University of Niš, Serbia

E-mail: [ljubicavelimirovic@yahoo.com](mailto:ljubicavelimirovic@yahoo.com)

<https://orcid.org/0000-0002-0317-4722>

The study of deformations dates from the ancient times and stems from purely practical problems. Deformations are in close connection with thin elastic shell and has a huge application from the mechanical point of view. In biology, the notion of deformation has also found his place. In geometry, the problem of deformations is covered by so-called the **surface bending theory**.

The surface bending theory considers the bending of surfaces, ie. the isometrical deformations, as well as, the infinitesimal bending of surfaces and presents one of the main parts of **global differential geometry**. In this talk, we want to introduce the manuscript of a book "Shape and Energies of Geometric Objects" whose authors are L. Kauffman, M. Najdanovic, S. Rancic and me. We consider shape of deformable objects at small, infinitesimal deformations. It is determined by their geometric curvature as well as curvature-based functionals which are said to be the energies of Geometric objects. Geometric objects that are discussed are curves surfaces and knots. To study energies and shape of curves and knots we use Willmore energy, total curvature, total torsion, total normalcy and Mbius energy. We expect that our investigations of energy and Willmore energy will eventually shed light on such phenomena.

## Structural Shape Analysis

**Nikola Velimirović<sup>1</sup>, Marija Najdanović<sup>2</sup>, Milan Zlatanović<sup>3</sup>,  
Andrija Zorić<sup>4</sup>**

<sup>1</sup>Department of Technical and Technological Sciences, State University of Novi  
Pazar, Serbia

E-mail: [nvelimirovic@np.ac.rs](mailto:nvelimirovic@np.ac.rs)

<https://orcid.org/0000-0002-6298-8216>

<sup>2</sup> Faculty of Sciences and Mathematics, University of Priština in Kosovska  
Mitrovica, Serbia

E-mail: [marija.najdanovic@pr.ac.rs](mailto:marija.najdanovic@pr.ac.rs)

<https://orcid.org/0000-0003-4149-9664>

<sup>3</sup>Faculty of Science and Mathematics, University of Niš, Serbia

E-mail: [milan.zlatanovic@pmf.edu.rs](mailto:milan.zlatanovic@pmf.edu.rs)

<https://orcid.org/0000-0002-0318-1092>

<sup>4</sup>Faculty of Civil Engineering and Architecture, University of Niš, Serbia

E-mail: [andrija.zoric@gaf.ni.ac.rs](mailto:andrija.zoric@gaf.ni.ac.rs)

<https://orcid.org/0000-0002-3107-9204>

Contemporary architectural and structural designs embrace irregular or freeform surfaces for their aesthetic appeal. The structural shape has important role in various aspects of design, analysis and construction. We have employed mathematical analysis of curvature and curvature based functions in order to understand surface characteristics. In civil engineering, adoption of the optimal structural shape is vital to guarantee structures fulfill essential performance criteria such as strength, stability, and functionality during the service life.

Keywords: structural shape, curvature, civil engineering

## **A generalization of a result of Minakshisundaram and Pleijel**

**Murali Vemuri**

Indian Institute of Technology (BHU), India  
E-mail: mkvemuri.mat@itbhu.ac.in

Minakshisundaram and Pleijel gave an asymptotic formula for the sum of squares of the pointwise values of the eigenfunctions of the Laplace-Beltrami operator on a compact Riemannian manifold, with eigenvalues less than a fixed number. Here, a generalization is given, where the pointwise values are replaced by the Fourier coefficients of a smooth measure supported on a compact submanifold.

## New invariants for conformal mappings of Riemannian spaces

Nenad Vesić<sup>1</sup>, Dušan Simjanović<sup>2</sup>

<sup>1</sup>Mathematical Institute of Serbian Academy of Sciences and Arts, Serbia

E-mail: vesic.specijalac@gmail.com

<https://orcid.org/0000-0001-7070-7325>

<sup>2</sup>Metropolitan University, Belgrade, Serbia

E-mail: dsimce@gmail.com

<https://orcid.org/0000-0002-1709-0765>

In this research, we are aimed to complete the study of invariants for conformal mappings of Riemannian spaces. Until now, the Weyl conformal tensor is the only invariant for this mapping which has been used. We obtained the invariant for conformal mapping  $f$  which is analogous to the Thomas projective parameter. After that, we obtained two invariants for conformal mappings with respect to transformation of Riemann-Christoffel curvature tensor of a Riemannian space. One of these two invariants is the Weyl conformal curvature tensor, but the second one is independent of this known invariant.

**Key words:** conformal mapping, basic invariants, weyl conformal tensor

**Math. Subj. Classification:** 53A55, 53B20

The first author is financially supported by Serbian Ministry of Science, Technological Development, and Patents, through the Mathematical Institute of Serbian Academy of Sciences and Arts

## On Yamada polynomial of spatial graphs and Jones polynomial of related links

Andrei Vesnin

Sobolev Institute of Mathematics and Tomsk State University, Russia

E-mail: andrei.vesnin@gmail.com

<https://orcid.org/0000-0001-7553-1269>

We will discuss invariants of spatial embeddings of graphs into a 3-sphere. Since any simple cycle of a graph is embedding as a knot, the theory of spatial graphs can be considered as a generalization of the knot theory. Two spatial graphs are said to be equivalent if there is an ambient isotopy of the 3-sphere which transforms one spatial graph to another. As well as knots and links, spatial graphs can be studied from their diagrams and Seifert surfaces. It was shown in [1] that an invariant link can be associated to an embedding a planar three-valent graph with at most six edges. The Yamada polynomial introduced in [2] is the most useful invariant of spatial graphs. A relation between the normalized Yamada polynomial of a spatial theta-graph and the Jones polynomial of the associated link was described in [3]. Let  $K_4$  be the complete graph with four vertices. We will present a relation between the normalized Yamada polynomials of a spatial  $K_4$ -graph and its spatial subgraphs with the Jones polynomial of the associated link, see [4] for details.

### References

- [1] L.H. Kauffman, J. Simons, K. Wolcott, P. Zhao, *Invariants of theta-curves and other graphs in 3-space*. Topology and Its Applications, 49 (1993), 193-216.
- [2] S. Yamada, *An invariant of spatial graphs*, Graph Theory, 13 (1989), 537-531.

- [3] Y. Huh, *Yamada polynomial and associated link of theta-curves*. Discrete Mathematics, 347:1 (2024), paper 113684.
- [4] O. Oshmarina, A. Vesnin, *Polynomials of complete spatial graphs and Jones polynomial of related links*. 2024, 28 pp. Preprint arXiv:2404.12264.

## Geodesics in Gromov-Hausdorff class

Anton Vihrov

Lomonosov Moscow State University, Moscow, Russian Federation

E-mail: Vihrov.09@gmail.com

<https://orcid.org/0009-0008-9852-2152>

The talk is dedicated to the proper class of all metric spaces considered up to isometry, equipped with the Gromov–Hausdorff distance. More precisely, we discuss the properties of geodesics in this Gromov–Hausdorff class.

The Gromov–Hausdorff distance measures the degree of the difference between two metric spaces. This distance was introduced by Gromov in 1981 and was defined as the smallest Hausdorff distance between isometric images of the considered spaces. Later, an equivalent definition of this distance was given using correspondences.

In Ivanov–Nikolaeva–Tuzhilin work an optimal correspondence between finite metric spaces was used to construct a geodesic between arbitrary compact metric spaces. Later on, the existence of optimal correspondence between compact metric spaces was proved, and as a consequence, a geodesic between these spaces generated by the optimal correspondence was constructed. Such geodesics are called *linear* ones. However, it is still unknown whether any pair of metric spaces at a finite distance from each other can be connected by some geodesic.

The example of two metric spaces without linear geodesic between them is trivial, e.g.  $[0,1]$  and  $(0,1)$ . Ghanaat found an example of two complete non- isometric metric spaces on zero distance, thus, they are not connected by linear geodesic. Hansen and the author found independently similar examples on non- zero distance. However, in the both examples, one can choose another metric spaces on zero distance from the initial ones, such that the latter spaces are connected by a linear geodesic. Later, the author found an example of two complete metric spaces such that no linear geodesics connect them and there are no metric spaces on zero distance. However, there exists a geodesic that are not linear. In the talk another method to construct a geodesic will be presented. It is called Hausdorff realization. In the case



two metric spaces have linear geodesic, then one can construct geodesic by means of Hausdorff realization. Nevertheless, there is an example of two metric spaces with no Hausdorff realization (Hansen). We will construct a geodesic in this case as well.

## Einstein Foliations

**Paweł Walczak**

Chair of Geometry, University of Łódź, Poland

E-mail: [pawel.walczak@wmii.uni.lodz.pl](mailto:pawel.walczak@wmii.uni.lodz.pl)

<https://orcid.org/0000-0001-8853-9622>

A Riemannian manifold is said to be *Einstein* whenever its Ricci curvature is constant. A foliation of a Riemannian manifold is *Einstein* whenever all its leaves are Einstein when equipped with induced Riemannian structures. Using Gauss and Codazzi equations, we shall provide conditions necessary and/or sufficient for this property and discuss the problem of existence of foliations of this sort. To simplify calculations and the formulas, we shall focus on the case when the ambient space is locally symmetric.

## On statistics which are almost sufficient from the viewpoint of the Fisher metrics

Kaori Yamaguchi

Ritsumeikan University, Japan  
E-mail: ra0097vv@ed.ritsumei.ac.jp

Given a statistical model, a statistic on the model is sufficient if the Fisher metric of the induced model coincides with the original Fisher metric, according to the definition by Ay-Jost-Lê-Schwachhöfer. We introduce and study its quantitative version: for  $0 < \delta \leq 1$ , we call a statistic  $\delta$ -almost sufficient if  $\delta^2 \mathbf{g}(v, v) \leq \mathbf{g}'(v, v)$  for every tangent vector  $v$  of the parameter space, where  $\mathbf{g}$  and  $\mathbf{g}'$  are the Fisher metric of the original and the induced model, respectively. By the monotonicity theorem, the Fisher metric of the induced model for such a statistic is bi-Lipschitz equivalent to the original one, which means that the information loss of the statistic is bounded in a quantitative way. We characterize such statistics in terms of the conditional probability or by the existence of a certain decomposition of the density function in a way similar to the characterizations of sufficient statistics due to Ay-Jost- Lê-Schwachhöfer and Fisher-Neyman.

## **Legendrian dual surfaces, lying in the 3-dimensional de Sitter space, of a spacelike curve in the 3-dimensional lightcone**

**Handan Yıldırım**

Istanbul University, Science Faculty, Mathematics Department, Istanbul  
Türkiye

E-mail: handanyildirim@istanbul.edu.tr

Legendrian dualities and the duality theorem were given in [1] for the pseudo-spheres in Lorentz-Minkowski space. Later, these Legendrian dualities and the duality theorem were extended in [2] for one-parameter families of pseudo-spheres in Lorentz-Minkowski space. In terms of two extended Legendrian dualities and the extended duality theorem in [2] and new extended Legendrian dualities in [3], an extended duality theorem was given in [3] and as an application of this theorem, Legendrian dual surfaces of a spacelike curve in the 3-dimensional lightcone were constructed in [3].

In this talk which is based on the joint work with Kentaro Saji given in [3], after constructing the Legendrian dual surface, lying in the 3-dimensional de Sitter space, of a spacelike curve in the 3-dimensional lightcone, the singularities of this surface are given.

### **Acknowledgements**

A part of [3] was conducted by the fruitful discussions of the speaker and Kentaro Saji during the speaker's participations to some scientific activities through the projects numbered 23429 and 33279 which were supported by the Scientific Research Projects Coordination Unit of Istanbul University. Moreover, a part of [3] was conducted during the speaker's visit to Mathematics Department of Kobe University in 2014. Furthermore, Kentaro Saji was partially supported by JSPS KAKENHI Grant Number 18K03301.

## References

- [1] S. Izumiya, *Legendrian dualities and spacelike hypersurfaces in the lightcone*, Moscow Mathematical Journal, **9**(2) (2009), 325-357.
- [2] S. Izumiya, H. Yildirim, *Extensions of the mandala of Legendrian dualities for pseudo-spheres in Lorentz-Minkowski space*, Topology and its Applications, **159**(2) (2012), 509-518.
- [3] K. Saji, H. Yildirim, *Legendrian dual surfaces of a spacelike curve in the 3-dimensional lightcone*, Journal of Geometry and Physics, **179** (2022) 104593, 1-21.

## Semisimple algebraic Nijenhuis operators on small dimensional Lie algebras

Ekaterina Zhikhareva

Lomonosov Moscow State University, Moscow, Russian Federation  
E-mail: marmeesha@yandex.ru

We consider a problem related to the Nijenhuis geometry.  
For arbitrary operator field  $L$  the formula

$$\mathcal{N}_A(\xi, \eta) = A[[A\xi, \eta]] + A[[\xi, A\eta]] - [[A\xi, A\eta]] - A^2[[\xi, \eta]]$$

defines  $(1, 2)$  type tensor field called *Nijenhuis torsion*. Here  $[[\ , \ ]]$  stands for standard commutator of vector fields. An operator field  $L$  is called *Nijenhuis operator* if its Nijenhuis torsion identically vanishes.

A natural class of such operators are left-invariant Nijenhuis operators  $A$  on a Lie group  $G$ . This means that  $A$  is invariant under the left action of the group on itself. In this case the tensor field  $A$  is completely defined by a linear operator  $L : g \rightarrow g$ , where  $g$  is the Lie algebra of  $G$ . Such operator  $L$  satisfies the identity

$$L[L\xi, \eta] + L[\xi, L\eta] - [L\xi, L\eta] - L^2[\xi, \eta] = 0,$$

where  $[\ , \ ]$  is the Lie algebra commutator. In this case the operator is called an *algebraic Nijenhuis operator*. Such operators appear in a study of completely integrable systems (both classic and quantum) and in many questions in geometry and mathematical physics. But we still have few examples of such operators. Our goal is to classify all algebraic Nijenhuis operators for small dimensional Lie algebras.

There is a theorem due to Y. Kosmann-Schwarzbach, F. Magri providing a characteristic property of such operators. This theorem says that real or complex Lie algebra  $g$  admits semisimple algebraic Nijenhuis operator if and only if there exists a basis  $\eta_1, \dots, \eta_n$  such that

$$[\eta_i, \eta_j] \in \langle \eta_i, \eta_j \rangle.$$

In other words any two vectors  $\eta_i, \eta_j$  generate a two-dimensional Lie subalgebra.

Applying this theorem we classified real and complex Lie algebras admitting algebraic Nijenhuis operators in dimensions three and four. For each such Lie algebra we describe all bases satisfying the above condition. An equally interesting question is to describe all algebraic Nijenhuis operators which are not equivalent by an automorphism of the Lie algebra. This problem is not solved yet, but we obtained the answer for some Lie algebras.

## **(Non) existence of Lagrangians in hyperkähler manifolds**

**Filip Živanović**

Simons Center for Geometry and Physics  
State University of New York, Stony Brook, NY 11794, United States  
E-mail: fzivanovic@scgp.stonybrook.edu

The purpose of the talk is to convey two phenomena, the existence and non-existence of Lagrangian submanifolds inside hyperkähler manifolds with holomorphic  $\mathbb{C}^*$ -actions. The first ones arise as the minima of the moment map of the corresponding  $S^1$ -actions, whereas the non-existence of the latter ones comes from the vanishing of a certain Floer-theoretic invariant called symplectic cohomology. We will discuss minimal resolutions of  $A_n$ -singularities as a basic example.



# Author Index

- Agapov  
    Sergei , 3  
Al Shaqsi  
    Khalifa , 5  
Aleksandar Lipkovski , 52  
Alekseevsky  
    Dmitri , 4  
Anton Savin , 86  
Arsenović  
    Miloš , 6  
Arutyunov  
    Andronick, 7  
Aydin  
    Muhittin Evren, 8  
  
Balashchenko  
    Vitaly , 9  
Baralić  
    Djordje , 11  
Bejan  
    Cornelia-Livia , 12  
Brozos-Vázquez  
    Miguel , 13  
  
Chen  
    Bang-Yen , 14  
Cvetković  
    Milica , 15  
Dekić  
    Andrijana, 16  
  
Deszcz  
    Ryszard , 17  
Dimitrijević  
    Ivan , 19  
Djorić  
    Miloš, 20  
Dragović  
    Vladimir, 22  
Dragovich  
    Branko, 21  
Dusek  
    Zdenek , 23  
  
Efremova  
    Lyudmila , 24  
Erken  
    İrem Küpeli, 25, 71  
  
Garcia-Rio  
    Eduardo , 27  
Gogić  
    Ilja , 28  
Guseva  
    Nadezda , 53  
  
Hinterleitner  
    Irena, 53  
Huijun  
    FAN , 26  
Huseini  
    Admir, 29

Ilyukhin	Velichka, 60
Daniil , 30	Minči´
Ilyutko	Svetislav, 61
Denis , 31	Mishchenko
Ivanov	Alexandr, 62
Stefan, 32	Mishra
	Mansi, 65
Jovanović	Mitrović
Božidar , 36	Ana , 15
Kauffman	Najdanović
Louis , 39	Marija , 66
Kocić	Najdanović
Djordje, 40	Marija, 99
Kudryavtseva	Neacşu
Elena , 41	Crina-Daniela, 67
	Nikolayevsky
Lazarov	Yuri, 69
Nenad Dj., 43	Nikonorov
Limonchenko	Yurii , 68
Ivan , 37	Nosovskiy
Lukić	Gleb , 70
Katarina, 45	
Makhrova	Ozkan
Elena , 46	Mustafa, 71
Maksimović	
Miroslav, 48	Patra
Manev	Dhriti Sundar , 72
Mancho, 49	Patrik Peška
Mateljević	P. , 53
Miodrag , 50	Pavlović
Matović	Milan, 73
Jelena Matović , 52	Petkov
Mikes	Alexander , 74
Josef, 34, 53	Petrović
Milenković	Miloš, 75
Vladislava , 56	Pitea
Millionshchikov	Ariana, 77
Dmitry , 57	Popelensky
Milousheva	Theodore, 78

Qing	van Coevering
Li, 44	Craig , 96
Radović	Velimirović
Ljiljana , 79	Ana, 97
Rančić	Ljubica, 98
Lidija , 80	Nikola, 99
Svetozar , 80	Vemuri
Rani	Murali, 100
Pooja, 81	Vesić
Rovenski	Nenad, 89, 101
Vladimir, 82, 84	Vesnín
Shubert	Andrei , 102
Anastasiia , 88	Vikhrov
Simjanović	Anton , 104
Dušan, 89, 101	Vilcu
Stanković	Gabriel-Eduard, 67
Mića S. , 90	Vukmirović
Stepanov	Srdjan , 36
Sergey E. , 91	Walczak
Stojanov	Paweł 106
Jelena, 92	Yıldırım
Stojanović	Handan , 108
Milica, 93	Yamaguchi
Sukilović	Kaori , 107
Tijana, 36, 73	Zhikhareva
Tuzhilin	Ekaterina , 110
Alexey, 94	Zivanović
Uljarević	Filip, 112
Igor, 95	Zlatanović
Vítková	Milan, 48, 97, 99
Lenka, 53	Zorić
	Andrija, 99